

On the equational theory of submodule lattices.

By Christian Herrmann

Equational problems for modular lattices have been studied for a long time, although available results have been established only under significant syntactical difficulties (see e.g. the papers of RALPH FREESE and ALEIT MITSCHKE in this volume). Furthermore, they have been more or less partial in nature. For lattices of submodules things become surprisingly easy, by simply making use of well known algebraic facts. As a by-product algebraic results can be extended by lattice theoretic methods. §1-3 are based on joint work of the author and A.HUHN; the results in §4 have been partially reported in [6].

§1 The two basic lemmas.

For  $p$  prime,  $n < \infty$ ,  $k \leq \infty$ , let  $L(p,k,n)$  be the lattice of subgroups of the  $n$ -th power of the cyclic group  $Z_{p^k}$  of order  $p^k$  or of the quasicyclic  $p$ -group  $Z_{p^\infty}$ . If  $M_R$  is a unitary  $R$ -module, then  $L(M_R)$  denotes the lattice of  $R$ -submodules of  $M_R$ . The lattice varieties generated by all normal subgroup lattices of groups or subgroup lattices of abelian groups or complemented modular lattices will be written as  $\mathcal{N}$  or  $\mathcal{R}$  or  $\mathcal{C}$ .  $\mathcal{K}$  shall denote the variety generated by  $\mathcal{N} \vee \mathcal{R}$ .

Lemma 1.  $L(M_R)$  is in the variety generated by all lattices  $L(p, k, n)$  where  $k < \infty$ ,  $p^k$  divides the characteristic of  $R$ , and  $n$  is less than or equal to the cardinality of a generating set of the  $P$ -module  $M_p$ , where  $P$  is the subring of  $R$  generated by the unit element.

Corollary 2.  $\mathcal{A}$  is generated by the finite primary lattices  $L(p, k, n)$  ( $p$  prime,  $k, n < \infty$ ).

Sketch of proof.  $L(M_R)$  is a sublattice of  $L(M_p)$  and  $L(M_p)$  is in the variety generated by the submodule lattices of its finitely generated submodules. By the Homomorphism Theorem these are sublattices of the  $L(P^n_p)$ . Now, if  $P$  is finite and  $|P| = p_1^{k_1} \dots p_m^{k_m}$ , then  $P \cong \prod_{i=1}^m (Z_{p_i^{k_i}})^n$  and  $L(P^n_p) \cong \prod_{i=1}^m L(p_i, k_i, n)$ . If, finally,  $P$  is isomorphic to the ring  $Z$  of integers, then we use the fact that a system of linear diophantine equations is solvable iff it is solvable in all  $Z_{p^k}$  and the following construction: To each lattice polynomial  $w$  attach a system  $\tilde{w}(x_i, y_i^j, \lambda_k)$  of linear equations in variables  $x_i, y_i^j, \lambda_k$  such that for any elements  $a_1, \dots, a_n, b_1^1, \dots, b_n^m$  of a  $R$ -module  $M_R$   $\langle a_1, \dots, a_n \rangle_R \in w(\langle b_1^1, \dots, b_n^1 \rangle_R, \dots, \langle b_1^m, \dots, b_n^m \rangle_R)$  holds iff the system  $\tilde{w}(a_i, b_i^j, \lambda_k)$  is solvable with values of the  $\lambda_k$  in  $R$ . This can be easily done by induction over the length of  $w$ . Hence, if all  $R$ -submodules of  $M_R$  are

generated by at most  $n$  elements, the inequality  $w \leq v$  is valid in  $L(M_R)$  if and only if, for any choice of constants  $a_i, b_i^j$  in  $M_R$ , the solvability of  $\tilde{w}(a_i, b_i^j, \lambda_k)$  implies the solvability of  $\tilde{v}(a_i, b_i^j, \mu_k)$  over  $R$ .

Lemma 3. If  $M_R$  is the  $\mathcal{F}$ -ultraproduct of the modules  $M_{iR_i}$ , then the  $\mathcal{F}$ -ultraproduct of the lattices  $L(M_{iR_i})$  is a sublattice of  $L(M_R)$ , containing all finitely generated  $R$ -submodules.

The proof is by the classical model theoretic method of correspondences between classes: consider the structures  $(M_R, L(M_R), \phi)$ ,  $\phi$  being the relation  $a \in U$  on  $M \times L(M_R)$  (c.f. MAKKAI, McNULTY [13]).

Corollary 4.  $\mathcal{C}$  is generated by subspace lattices of finite projective geometries over prime fields and arbitrary nondesarguesian planes.

## §2 Lattices generated by a frame.

In [9] A.HUHN introduced the concept of an  $n$ -diamond in a lattice: a sequence  $a_0, \dots, a_n$  of elements such that any  $n$ -element subset is independent in the interval  $[\prod_{i=0}^n a_i, \sum_{i=0}^n a_i]$ . It is called a frame in  $L$ , if  $\prod a_i = 0_L$

and  $\sum a_i = 1_L$ . If  $L$  is modular, a frame in the usual sense can be derived and vice versa.

Theorem 5. For  $n \geq 3$  there is a complete list of all subdirectly irreducible lattices in  $\mathcal{N}$  which are generated by an  $n$ -diamond:

the rational projective geometry  $L(Q^n_Q)$ ;

the lattices  $L(p, k, n)$ , where  $p$  is prime and  $k \leq \infty$ ;

the duals of the  $L(p, \infty, n)$ , where  $p$  is prime.

The generating  $n$ -diamond is given, up to automorphism, by the submodules  $e_0 = (x, \dots, x)$ ,  $e_i = (0, \dots, 0, x, 0, \dots, 0)$ , with  $x$  in the  $i$ -th entry, for  $i = 1, \dots, n$ .

The following notation has been used: For  $k_1, \dots, k_n$  in  $R$  and variables  $x_1, \dots, x_n$  not necessarily distinct we have  $(k_1 x_1, \dots, k_n x_n) = \{(k_1 a_1, \dots, k_n a_n) \mid a_i \in M \text{ and } x_i = x_j \Rightarrow a_i = a_j\}$ , a submodule of  $M_R^n$ .

The proof consists of the following main steps:

- 1) Reduction to  $\mathcal{R}$ : If the lattice of normal subgroups of  $G$  contains an  $n$ -frame ( $n \geq 3$ ), then  $G$  is abelian.
- 2) If  $L$  is a sublattice of any  $L(p, k, n)$  ( $k < \infty$ ) generated by an  $n$ -diamond, then  $L$  is a  $\{0, \dots, n\}$ -subdirect product (in the sense of WILLE [18]) of lattices  $L(p_i, k_i, n)$ , each generated by the diamond  $e_0, \dots, e_n$ .

- 3) By Corollary 2 the lattice  $FR(P_n)$  freely generated in  $\mathcal{A}$  by an  $n$ -diamond is a  $\{0, \dots, n\}$ -subdirect product of the lattices  $L(p, k, n)$  ( $p$  prime,  $k < \infty$ ) with generators  $e_0, \dots, e_n$ .
- 4) Any subdirectly irreducible lattice in  $\mathcal{A}$  which is generated by an  $n$ -diamond is, using the Lemma in JÓNSSON [12], a homomorphic image of the sublattice  $M$  generated by  $[e_i]_{\mathcal{F}}$  ( $i=0, \dots, n$ ) in  $FR(P_n)/\mathcal{F}$  for a suitable ultrafilter  $\mathcal{F}$  on the set  $\{p^k \mid p \text{ prime, } k < \infty\}$ . Now, if  $p \rightarrow \infty$  in  $\mathcal{F}$ , then  $M \cong L(Q^n_Q)$  is proved with the method of Lemma 3. If, on the other hand,  $k \rightarrow \infty$  for fixed  $p$ , then  $M$  is a subdirect product of  $L(p, \infty, n)$  and its dual. But the only nontrivial homomorphic image of these lattices is  $L(p, 1, n)$ . [Ralph Freese observed that the image is  \$L\(Q^n\_Q\)\$ .](#)

Corollary 6. The subdirectly irreducible lattices in  $\mathcal{C}$  generated by an  $n$ -diamond are, for  $n \geq 3$ , exactly those in the above list and, for  $n=3$ , those in the above list as well as nondesarguesian planes generated by four points.

§3 Applications to equational classes

In order to apply Theorem 5 we need polynomials

$d_i(x_0, \dots, x_3)$  ( $i=0, \dots, 3$ ) such that in any modular lattice the following holds: For any choice of  $x_0, \dots, x_3$  the  $d_i(x_0, \dots, x_3)$  ( $i=0, \dots, 3$ ) are equal to each other or form a **3**-diamond; if  $x_0, \dots, x_3$  is a **3**-diamond, then  $x_i = d_i(x_0, \dots, x_3)$  ( $i=0, \dots, 3$ ). Such polynomials are defined in A.HUHN [9] :

$$d_0(x_0, \dots, x_3) := \prod_{i=1}^3 b_i(x_0, \dots, x_3) + a_0(x_0, \dots, x_3)$$

$$d_i(x_0, \dots, x_3) := \prod_{j=1, j \neq i}^3 b_j(x_0, \dots, x_3) \quad \text{for } i=1, 2, 3$$

where  $a_0(x_0, \dots, x_3) = x_0 \cdot \sum_{i=1}^3 x_i$ ,  $a_i(x_0, \dots, x_3) = \sum_{j=1, j \neq i}^3 x_j$ ,

$$v(x_0, \dots, x_3) = \prod_{i=1}^3 a_i(x_0, \dots, x_3) + a_0(x_0, \dots, x_3)$$

$$b_i(x_0, \dots, x_3) = a_i(x_0, \dots, x_3) \cdot v(x_0, \dots, x_3) + \sum_{j=1}^3 x_0 \cdot \sum_{k=1, k=j}^3 x_k$$

for  $i=1, 2, 3$ .

Now we can define inductively  $w_0(x_0, \dots, x_3) = d_3(x_0, \dots, x_3)$

$$w_{n+1}(x_0, \dots, x_3) = \left\{ \left[ w_n(x_0, \dots, x_3) + d_0(x_0, \dots, x_3) \cdot d_2(x_0, \dots, x_3) \right] \cdot \left[ d_1(x_0, \dots, x_3) + d_2(x_0, \dots, x_3) \right] + d_0(x_0, \dots, x_3) \right\} \cdot \left[ d_2(x_0, \dots, x_3) + d_3(x_0, \dots, x_3) \right]$$

Lemma 7.  $w_n(e_0, \dots, e_3) = (0, nx, x)$  in any module  $M_R^3$ .

Theorem 8. The lattice identity  $w_n(x_0, \dots, x_3) = d_3(x_0, \dots, x_3)$  is valid in  $L(M_R)$  if and only if the greatest common divisor of the additive orders of any three weakly independent elements of  $M$  divides  $n$ .

Theorem 9. Each lattice  $L(p, k, n)$  ( $n \geq 3$ ,  $k < \infty$ ) is splitting in  $\mathcal{H}$ . For  $k > 1$  splitting universal disjunctions are "length  $\leq 3$ " or  $d_3(x_0, \dots, x_3) \cdot w_p^k(x_0, \dots, x_3) \leq d_3(x_0, \dots, x_3) \cdot w_p^{k-1}(x_0, \dots, x_3)$ , and for  $k=1$ , they are "length  $\leq 3$  and  $L(p, k, n)$  not order embeddable" or  $d_3(x_0, \dots, x_3) \cdot w_p(x_0, \dots, x_3) \leq x$ . We remark that  $L(p, 1, n)$  is neither projective nor finitely projected nor bounded epimorphic image of a free lattice. [In contrast, Ralph Freese showed that  \$L\(p, 1, n\)\$  is projective in the class of modular lattices](#)

Theorem 10. If  $\mathcal{L}$  is any class of lattices contained in  $\mathcal{H}$  and containing all sublattices of lattices  $L(V_K)$  where  $V_K$  is any five dimensional vector space over a field of characteristic zero, then  $\mathcal{L}$  cannot be defined by a finite set of first order axioms.

The proof is immediate by the following Lemma 11 and the fact that a nontrivial ultraproduct of  $L_{p,q}$ 's is embeddable in a  $L(V_K)$ , (see [5]).

Lemma 11. There is an identity valid in  $\mathcal{H}$  which does not hold, for  $p \neq q$ , in the Arguesian lattice  $L_{p,q} = [0, a] \cup [b, 1]$  with  $b \not\leq a$ ,  $[0, a] \cong L(p, 1, 3)$ , and  $[b, 1] \cong L(q, 1, 3)$  - cf. JOHNSON [11].

#### §4 Lattices with four generators.

In [1] the authors asked for a complete list of subdirectly irreducible modular lattices with four generators; the solution is still distant.

Lemma 12. Any lattice listed in Theorem 5 is generated by four elements.

Furthermore, with the methods of [7] it is possible to construct from a sufficiently large partial sublattice of a lattice  $L(p,2,3)$  a nondesarguesian uniform Hjelmslevplane with four generators.

The systems of generators and a partial converse of the lemma stem from the work of GELFAND and PONOMAREV [3] on linear spaces with four subspaces.

If  $V$  is a linear space with subspaces  $V_1, \dots, V_4$ , then  $(V, V_1, \dots, V_4)$  is called a quadruple. It is called indecomposable, if there is no nontrivial complementary pair  $A, B$  of subspaces such that  $V_i = A \cap V_i + B \cap V_i$  for  $i=1, \dots, 4$ .

Lemma 13. If  $L = \langle v_1, \dots, v_4 \rangle$  is subdirectly irreducible and can be embedded in the subspace lattice of a linear space of finite dimension, then there is an indecomposable quadruple  $(V, V_1, \dots, V_4)$  (with  $V$  of finite dimension over an algebraically closed field  $F$ ) and an isomorphism of  $L$  onto the sublattice  $\langle V_1, \dots, V_4 \rangle$  of  $L(V_F)$  mapping  $v_i$  onto  $V_i$  for  $i=1, \dots, 4$ .



Theorem 14. (GELFAND, PONOMAREV [3]) The indecomposable quadruples of finite dimensions over an algebraically closed field  $F$  are given (up to isomorphism, permutation, and duality) by the following list:

- 1)  $(F^{2n}, (x^n, 0^n), (0^n, x^n), (x^n, x^n), (x^1, y^{n-1}, \lambda x^1, \lambda y^{n-1} + x^1))$   
with  $\lambda \in F - \{0, 1\}$ ,
- 2) a)  $(F^{2n+1}, (x^{n+1}, 0^n), (0^{n+1}, x^n), (x^1, y^n, y^n), (x^n, 0^1, x^n))$  ,  
b)  $(F^{2n}, (x^n, 0^n), (0^n, x^n), (x^1, y^{n-1}, y^{n-1}, 0^1), (x^n, x^n))$  ,
- 3) a)  $(F^{2n+1}, (x^n, 0^{n+1}), (0^n, x^{n+1}), (x^n, x^n, 0^1), (x^n, 0^1, x^n))$  ,  
b)  $(F^{2n}, (x^n, 0^n), (0^n, x^n), (0^1, x^{n-1}, x^{n-1}, 0^1), (x^n, x^n))$  ,
- 4)  $(F^{2n+1}, (x^n, 0^{n+1}), (0^n, x^n, 0^1), (0, x_1, \dots, x_{n-1}, x_1, \dots, x_n, x_n),$   
 $(x_1, \dots, x_n, 0, x_1, \dots, x_n))$  .

Let  $S(n, 4)$  be the lattice of fig.1 and  $FM(J_1^4)$  the modular lattice freely generated by  $J_1^4$  (cf. fig.2 and [1]) .

Lemma 15.  $V_1, \dots, V_4 \subseteq L(V_F)$  is in cases 1)-4) isomorphic to  $M_4, S(m, 4), L(P_P^m)$  where  $m=2n+1$  or  $m=2n$  and  $P$  the prime field of  $F$ , respectively.

Theorem 16. If  $L$  is subdirectly irreducible in  $\mathcal{C}$  and generated by four elements, then  $L$  is isomorphic either to a nondegenerate plane or one of the following lattices:

$M_4, S(m, 4), L(P_P^m)$   $P$  a prime field,  $FM(J_1^4)$  or its dual .

Sketch of proof. Similarly as in the proof of Theorem 3 we

have to study ultraproducts of lattices listed in Lemma 15. As there is only a finite number of types, we may assume that all components are of the same type. In case 1) there is nothing to do; in 2) the untraproduct is again of breadth two, hence we may use the result of FREESE [2] that any subdirectly irreducible breadth two modular lattice with four generators is  $S(m,4)$ ,  $FM(J_1^4)$  or its dual. But the sublattice  $M$  generated by  $v_1, \dots, v_4$  in the ultraproduct can be visualized and decomposed in a straightforward manner.

3), 4): If  $m$  is fixed, then  $M \cong L(P^m_P)$  follows trivially.

If  $m \rightarrow \infty$ , then we consider structures

$(P^m_P, L(P^m_P), v_1, \dots, v_4, \Phi, I, J, K, \kappa, \varphi, 1_I, 1_J, \infty_I, \infty_J)$  such that:  $(P^m_P, v_1, \dots, v_4)$  is the given quadruple;  $\Phi$  is defined as in Lemma 3;  $I = \{1, \dots, n\}$ ,  $J = \{n+1, \dots, 2n+1\}$  in case 3a),  $J = \{n+1, \dots, 2n\}$  in cases 3b) and 4),  $K = \emptyset$  in cases 3 a) and b), and  $K = \{2n+1\}$  in cases 4);  $I$  and  $J$  are equipped with the partial operation of taking the successor;  $\varphi$  is the mapping from  $I$  onto  $J$  with  $\varphi(i) = i+n$ ;  $\kappa$  is the mapping from  $P^m_{I \cup J \cup K}$  into  $P$  such that  $\kappa(a, i)$  is the  $i$ th coordinate of  $a$ ;  $1_I = 1$ ,  $1_J = n+1$ ,  $\infty_I = n$ ,  $\infty_J = 2n+1$  in case 3a), and  $\infty_J = 2n$  in cases 3b) and 4).

In any of the cases 3a), 3b), and 4) we have formulas  $\alpha_1, \dots, \alpha_4$  in the first order language of these structures such that for any  $m$  and  $P$   $v_i = \{x \mid x \in P^m \text{ and } \alpha_i(x)\}$  holds for  $i=1, \dots, 4$ .

Now, in the ultraproduct we have a vector space  $V \cong F^{I \cup J \cup K}$  and

$v_i = \{x \mid x \in V \text{ and } \alpha_i(x)\}$  is valid as well.

Let  $I_1$  and  $I_\infty$  be the subalgebras of  $I \cup J$  generated by  $\{1_I, 1_J\}$  and  $\{\omega_I, \omega_J\}$  resp., and  $I_* = (I \cup J) - (I_1 \cup I_\infty)$ . Define  $A_\gamma = \{f \mid f \in V \text{ and } (f, i) = 0 \text{ for all } i \in I_\gamma\}$ , a subspace of  $V$ , for  $\gamma \in \{1, *\}$  or  $\gamma = \infty$  and case 3a, b) ;  $A_\infty = \{f \mid f \in V \text{ and } (f, i) = 0 \text{ for all } i \in I_\infty \cup K\}$  in case 4) .

Then  $A_1, A_*, A_\infty$  yield a direct decomposition of the quadruple  $(V, V_1, \dots, v_4)$  into three quadruples  $(A_\gamma, v_1^\gamma, \dots, v_4^\gamma)$  ( $\gamma = 1, *, \infty$ ), thus a subdirect decomposition of  $M$  into three factors.

But for  $\gamma = 1, *$  or  $\gamma = \infty$  and case 3a, b)  $v_1^\gamma, \dots, v_4^\gamma$  together with  $0$  and  $V$  form a partial lattice  $J_1^4$  ; hence they generate a lattice  $FM(J_1^4)$  (cf. [1]). In case 4) from  $v_1^\infty, \dots, v_4^\infty$  we get a partial lattice  $J_3^4$  (fig.3) which generates a third subdirect power of  $FM(J_1^4)$  (see [14]). In any case,  $M$  is a finite subdirect power of  $FM(J_1^4)$  and the only subdirectly irreducible epimorphic images of  $M$  are  $M_4$  and  $FM(J_1^4)$  .

### §5 Word problems.

HUTCHINSON [10] proved that in a quasivariety  $\mathcal{Q}$  of modular lattices such that  $L(R^{\omega}_R) \in \mathcal{Q}$  for a nontrivial ring  $R$  there is a finitely presented lattice with seven generators which has an unsolvable word problem (cf. FREESE [2], too). The attempts on the word problem for free modular lattices  $FM(n)$  by SCHÜTZENBERGER [16] and GLUHOV [4] may be regarded as unsuccessful (cf. WHITMAN [17] and HERRMANN [8]). The following solvability results do, however, hold. Here

${}_m\mathcal{M}^n$  denotes the class of all modular lattices of primitive length  $\leq n$  and primitive breadth  $\leq m$ .

Theorem 17. (FREESE 2) In  ${}_2\mathcal{M}$  the word problem in four generators is solvable.

Theorem 18. [7] For  $n \leq 6$  and  $m \leq 3$  or  $n < \infty$  and  $m \leq 2$  the word problem in  ${}_m\mathcal{M}^n$  is solvable.

Theorem 19. In  $\mathcal{C}$  the word problem in four generators is solvable.


Theorem 20. The word problems for the free lattices  $FC(n)$  and  $FA(n)$  are solvable.

Proof. By Theorem 16 any four generated lattice in  $\mathcal{C}$  is embeddable in a complemented modular lattice. Hence a Horn formula in four variables is valid in  $\mathcal{C}$  if and only if it can be derived from the finite set of axioms of complemented modular lattices by a calculus of first order logic. On the other hand the four variable  $\mathcal{C}$  Horn formulas not valid in  $\mathcal{C}$  are enumerable by Theorem 16, too. universal

Theorem 20 is an immediate consequence of Corollaries 2 and 4 and the fact that  $\mathcal{C}$  and  $\mathcal{A}$  can be defined by enumerable sets of identities. For  $\mathcal{C}$  these are just the identities derivable from the axioms of a complemented modular lattice; for  $\mathcal{A}$  this

follows by the result of SCHEIN[15] that the class of lattices embeddable in subgroup lattices of abelian groups can be recursively defined.


Fig.1



$$[0,1] = S(8,4)$$

$$[0,a] = S(7,4)$$

Fig.2




$$1 = a+b = a+c = a+d = b+d = c+d$$

$$0 = ab = ac = ad = bc = bd = cd$$

$J_1^4$

Fig.3



$$1 = a+c = a+d = b+d$$

$$0 = ab = ac = ad = bc = bd = cd$$

$J_3^4$

## Literature.

- 1 A.Day,C.Herrmann,R.Wille, On modular lattices with four generators, Algebra Universalis 2(1972), 317-323
- 2 R.Freese, Breadth two modular lattices, this volume
- 3 I.M.Gelfand,V.A.Ponomorev, Problems of linear algebra and classification of quadruples of subspaces in a finite-dimensional vector space, Coll.Math.J.Bolyai, 5,Hilbert Space Operators, Tihany 1970
- 4 M.M.Gluhov, Algorithmic solvability of the word problem for completely free modular lattices, Sibirsk.Math.Ž. 5(1964), 1027-1034
- 5 C.Herrmann,W.Poguntke, Axiomatic classes of lattices of normal subgroups, mimeographed notes, Darmstadt 1972
- 6 C.Herrmann,C.M.Ringel,R.Wille, On modular lattices with four generators, to appear in Notices AMS 1973
- 7 C.Herrmann, Modulare Verbände von Länge $\leq 6$ , This volume
- 8 C.Herrmann, Concerning M.M.Gluhovs paper on the word problem for free modular lattices, to appear
- 9 A.Huhn, On a problem of G.Grätzer, to appear
- 10 G.Hutchinson, Recursively unsolvable word problems of modular lattices and diagram-chasing, to appear
- 11 B.Jónsson, Algebras whose congruence lattices are distributive, Math.Scand. 21(1967),110-121
- 12 B.Jónsson, Modular lattices and Desargues' theorem, Math.Scand. 2(1954), 295-314
- 13 M.Makkai,G.McNulty, On axioms for lattices of submodules, mimeographed notes, Winnipeg 1973
- 14 G.Sauer,W.Seibert,R.Wille, On free modular lattices over partial lattices with four generators, this volume
- 15 B.M.Schein, Relation algebras and function semigroups, Semigroup Forum 1(1970), 1-62
- 16 M.P.Schützenberger, Le probleme des mots dans les treillis modulaires libres, C.R.Acad.Sci.Paris vol.237(1953),507-8
- 17 W.Whitman, Status of word problems for lattices, in Lattice Theory II, Proc.Symp.pure Math.AMS II(1961),17-21
- 18 R.Wille, Subdirekte Produkte und konjunkte Summen, J.reine und angew.Math. 239(1970), 333-8