

UNIT-REGULARITY AND REPRESENTABILITY FOR SEMIARTINIAN *-REGULAR RINGS. ERRATUM

CHRISTIAN HERRMANN

ABSTRACT. We show that any semiartinian *-regular ring R is unit-regular; if, in addition, R is subdirectly irreducible then it admits a representation within some inner product space.

0. ERRATUM

There is no proof of Thm. 7. since the quoted Fact 5 is incorrect.

1. INTRODUCTION

The motivating examples of *-regular rings, due to Murray and von Neumann, were the *-rings of unbounded operators affiliated with finite von Neumann algebra factors; to be subsumed, later, as *-rings of quotients of finite Rickart C^* -algebras. All the latter have been shown to be *-regular and unit-regular (Handelman [5]). Representations of these as *-rings of endomorphisms of suitable inner product spaces have been obtained first, in the von Neumann case, by Luca Giudici (cf. [6]), in general in joint work with Marina Semenova [9]. The existence of such representations implies direct finiteness [7]. In the present note we show that every semiartinian *-regular ring is unit-regular and a subdirect product of representables. This might be a contribution to the question, asked by Handelman (cf. [3, Problem 48]), whether all *-regular rings are unit-regular. We rely heavily on the result of Baccella and Spinosa [1] that a semiartinian regular ring is unit-regular provided that all its homomorphic images are directly finite. Also, we rely on the theory of representations of *-regular rings developed by Florence Micol [12] (cf. [9, 10]). Thanks are due to the referee for a timely, concise, and helpful report.

2. PRELIMINARIES: REGULAR AND *-REGULAR RINGS

We refer to Berberian [2] and Goodearl [3]. Unless stated otherwise, rings will be associative, with unit 1 as constant. A (von Neumann)

2000 *Mathematics Subject Classification.* Primary:16E50, 16W10.

Key words and phrases. *-regular ring, representable, unit-regular.

regular ring R is such that for each $a \in R$ there is $x \in R$ such that $axa = a$; equivalently, every right (left) principal ideal is generated by an idempotent.

The *socle* $Soc(M)$ of a right R -module is the sum of all minimal submodules. For a ring R define its *Loewy series* of right ideals $L_\alpha(R)$ by $L_0(R) = 0$, $L_{\alpha+1} = Soc(R/L_\alpha(R))$, and $L_\alpha(R) = \bigcup_{\beta < \alpha} L_\beta(R)$ is α is a limit ordinal. R has *Loewy length* α if $R = L_\alpha(R)$ with α minimal, provided that such exists. A ring R with unit is (right) *semiartinian* if R/M has nonzero socle for each right ideal of R ; equivalently, R has Loewy length α for some α - which must be of the form $\xi + 1$ since R has unit 1. If R is regular, then the $L_\alpha(R)$ are, moreover, ideals since left and right socle of a regular ring coincide [11].

A ring R is *directly finite* if $xy = 1$ implies $yx = 1$ for all $x, y \in R$. A ring R is *unit-regular* if for any $a \in R$ there is a unit u of R such that $aua = a$. Unit-regular rings are directly finite, in particular. The crucial fact to be used, here, is the following result of Baccella and Spinosa [1].

Theorem 1. *A semiartinian regular ring is unit-regular provided all its homomorphic images are directly finite.*

A **-ring* is a ring R endowed with an involution $r \mapsto r^*$. Such R is **-regular* if it is regular and $rr^* = 0$ only for $r = 0$. A *projection* is an idempotent e such that $e = e^*$; we write $e \in P(I)$ if $e \in I$. A **-ring* is **-regular* if and only if for any $a \in R$ there is a projection e with $aR = eR$; such e is unique and obtained as aa^+ where a^+ is the pseudo-inverse of a . In particular, for **-regular* R , each ideal I is a **-ideal*, that is, closed under the involution. Thus, R/I is a **-ring* with involution $a + I \mapsto a^* + I$ and a homomorphic image of the **-ring* R . In particular, R/I is regular; and **-regular* since $aa^+ + I$ is a projection generating $(a + I)(R/I)$.

If R is a **-regular* ring and $e \in P(R)$ then the *corner* eRe is a **-regular* ring with unit e , operations inherited from R , otherwise. For a **-regular* ring, $P(R)$ is a modular lattice, with partial order given by $e \leq f \Leftrightarrow fe = e$, which is isomorphic to the lattice $L(R)$ of principal right ideals of R via $e \mapsto eR$. In particular, eRe is artinian if and only if e is contained in the sum of finitely many minimal right ideals.

A **-ring* is *subdirectly irreducible* if it has a unique minimal ideal, denoted by $M(R)$. Observe that $Soc(R) \neq 0$ implies $M(R) \subseteq Soc(R)$ since $Soc(R)$ is an ideal. For the following see Lemma 2 and Theorem 3 in [8].

Fact 2. *If R is a subdirectly irreducible *-regular ring then eRe is simple for all $e \in P(M(R))$ and R a homomorphic image of a *-regular sub-*-ring of some ultraproduct of the eRe , $e \in P(M(R))$.*

3. PRELIMINARIES: REPRESENTATIONS

We refer to Gross [4] and Sections 1 of [9], 2–4 of [10]. By an *inner product space* V_F we will mean a right vector space (also denoted by V_F) over a division *-ring F , endowed with a sesqui-linear form $\langle \cdot | \cdot \rangle$ which is *anisotropic* ($\langle v | v \rangle = 0$ only for $v = 0$) and *orthosymmetric*, that is, $\langle v | w \rangle = 0$ if and only if $\langle w | v \rangle = 0$. Let $\text{End}^*(V_F)$ denote the *-ring consisting of those endomorphisms φ of the vector space V_F which have an adjoint φ^* w.r.t. $\langle \cdot | \cdot \rangle$.

A *representation* of a *-ring R within V_F is an embedding of R into $\text{End}^*(V)$. R is *representable* if such exists. The following is well known, cf. [11, Chapter IV.12]

Fact 3. *Each simple artinian *-regular ring is representable.*

The following two facts are consequences of Propositions 13 and 25 in [9] (cf. Micol [12, Corollary 3.9]) and, respectively, [7, Theorem 3.1] (cf. [8, Theorem 4]).

Fact 4. *A *-regular ring is representable provided it is a homomorphic image of a *-regular sub-*-ring of an ultraproduct of representable *-regular rings.*

Fact 5. *Every representable *-regular ring is directly finite.*

4. MAIN RESULTS

Theorem 6. *If R is a subdirectly irreducible *-regular ring such that $\text{Soc}(R) \neq 0$, then $\text{Soc}(R) = M(R)$, each eRe with $e \in P(M(R))$ is artinian, and R is representable.*

Proof. Consider a minimal right ideal aR . As R is subdirectly irreducible, $M(R)$ is contained in the ideal generated by a ; that is, for any $0 \neq e \in P(M(R))$ one has $e = \sum_i r_i a s_i$ for suitable $r_i, s_i \in R$, $r_i a s_i \neq 0$. By minimality of aR , one has $a s_i R = aR$ and $r_i a s_i R = r_i a R$ is minimal, too. Indeed, $x \mapsto r_i x$ is an R -linear map of aR onto $r_i a R \neq 0$. Thus, $e \in \sum_i r_i a R$ means that eRe is artinian. By Facts 3, 2, and 4, R is representable.

It remains to show that $\text{Soc}(R) \subseteq M(R)$. Recall that the congruence lattice of $L(R)$ is isomorphic to the ideal lattice of R ([13, Theorem 4.3] with an isomorphism $\theta \mapsto I$ such that $aR/0 \in \theta$ if and only if $a \in I$). In particular, since R is subdirectly irreducible so is $L(R)$.

Choose $e \in M(R)$ with eR minimal. Then for each minimal aR one has $eR/0$ in the lattice congruence θ generated by $aR/0$. Since both quotients are prime, by modularity this means that they are projective to each other. Thus, $aR/0$ is in the lattice congruence generated by $eR/0$ whence a is in the ideal generated by e , that is, in $M(R)$. \square

Theorem 7. *Every semiartinian $*$ -regular ring R is unit-regular and a subdirect product of representable homomorphic images.*

Proof. Consider an ideal I of R . Then $I = \bigcap_{x \in X} I_x$ with completely meet irreducible I_x , that is, subdirectly irreducible R/I_x . Since R is semiartinian one has $\text{Soc}(R/I_x) \neq 0$, whence R/I_x is representable by Theorem 6 and directly finite by Fact 5. Then R/I is directly finite, too, being a subdirect product of the R/I_x . By Theorem 1 it follows that R is unit-regular. \square

5. EXAMPLES

It appears that semiartinian $*$ -regular rings form a very special subclass of the class of unit-regular $*$ -regular rings, even within the class of those which are subdirect products of representables. E.g. the $*$ -ring of unbounded operators affiliated to the hyperfinite von Neumann algebra factor is representable, unit-regular, and $*$ -regular with zero socle. On the other hand, due to the following, for every simple artinian $*$ -regular ring R and any natural number $n > 0$ there is a semiartinian $*$ -regular ring having ideal lattice an n -element chain and R as a homomorphic image.

Proposition 8. *Every representable $*$ -regular ring R embeds into some subdirectly irreducible representable $*$ -regular ring \hat{R} such that $R \cong \hat{R}/M(\hat{R})$. In particular, \hat{R} is semiartinian if and only if so is R .*

The proof needs some preparation. Call a representation $\iota : R \rightarrow \text{End}^*(V_F)$ *large* if for all $a, b \in R$ with $\text{im } \iota(b) \subseteq \text{im } \iota(a)$ and finite $\dim(\text{im } \iota(a)/\text{im } \iota(b))_F$ one has $\text{im } \iota(a) = \text{im } \iota(b)$.

Lemma 9. *Any representable $*$ -regular ring admits some large representation.*

Proof. Inner product spaces can be considered as 2-sorted structures V_F with sorts V and F . In particular, the class of inner product spaces is closed under formation of ultraproducts. Representations of $*$ -rings R can be viewed as R - F -bimodules ${}_R V_F$, that is as 3-sorted structures, with R acting faithfully on V . It is easily verified that the class of representations of $*$ -rings is closed under ultraproducts cf. [9, Proposition 13].

Now, given a representation η of R in W_F , form an ultrapower ι , that is ${}_S V_{F'}$, such that $\dim F'_F$ is infinite (recall that F' is an ultrapower of F). Observe that $\mathbf{End}^*(V_{F'})$ is a sub- $*$ -ring of $\mathbf{End}^*(V_F)$ and $\dim(U/W)_F$ is infinite for any subspaces $U \supseteq W$ of $V_{F'}$. Also, S is an ultrapower of R with canonical embedding $\varepsilon : R \rightarrow S$. Thus, $\varepsilon \circ \iota$ is a large representation of R in V_F . \square

Proof. of Proposition 8. In view of Lemma 9 we may assume a large representation ι of R in V_F . Identifying R via ι with its image, we have R a $*$ -regular sub- $*$ -ring of $\mathbf{End}^*(V_F)$. Let I denote the set of all $\varphi \in \mathbf{End}(V_F)$ such that $\dim(\text{im } \varphi)_F$ is finite. According to Micol [12, Proposition 3.12] (cf. Propositions 4.4 (i),(iii) and 4.5 in [10]) $R + I$ is a $*$ -regular sub- $*$ -ring of $\mathbf{End}^*(V_F)$, with unique minimal ideal I . By Theorem 6 one has $I = \text{Soc}(R + I)$. Moreover, $R \cap I = \{0\}$ since the representation ι of R in V_F is large. Hence, $R \cong (R + I)/I$. \square

REFERENCES

- [1] Baccella, G., Spinoso, L.: K_0 of semiartinian von Neumann regular rings. *Direct finiteness versus unit-regularity*. *Algebr. Represent. Theory* 20 (2017), 1189–1213
- [2] S. K. Berberian, *Baer *-rings*, Springer, Grundlehren 195, Berlin 1972.
- [3] K. R. Goodearl, *Von Neumann Regular Rings*, second edition, Krieger, Malabar, 1991.
- [4] H. Gross, *Quadratic Forms in Infinite Dimensional Vector spaces*, Birkhuser, Basel, 1979.
- [5] D. Handelman, *Finite Rickart C^* -algebras and their properties*, in *Studies in analysis*, ed. G.-C. Rota, pp. 171–196, *Adv. in Math. Suppl. Stud.*, 4, Academic Press, New York-London, 1979.
- [6] Herrmann, C.: On the equational theory of projection lattices of finite von Neumann factors. *The Journal of Symbolic Logic*, Vol. 75, No. 3 (2010), 1102–1110
- [7] Herrmann, C., Direct finiteness of representable regular -rings. *Algebra Universalis* 80 (2019), no. 1, Art. 3, 5 pp. <http://arxiv.org/abs/1904.04505>
- [8] Herrmann, C., Varieties of $*$ -regular rings. <http://arxiv.org/abs/1904.04505>
- [9] Herrmann, C., Semenova, M. V., Rings of quotients of finite AW-algebras. *Representation and algebraic approximation*. (Russian) *Algebra Logika* 53 (2014), no. 4, 466–504, 550–551; translation in *Algebra Logic* 53 (2014), no. 4, 298–322
- [10] Herrmann, C., Semenova, M., Linear representations of regular rings and complemented modular lattices with involution. *Acta Sci. Math. (Szeged)* 82 (2016), no. 3–4, 395–442.
- [11] N. Jacobson, *Structure of Rings*, AMS Col. Publ. XXXVII, Amer. Math. Soc., Providence, RI, 1956
- [12] F. Micol, *On representability of $*$ -regular rings and modular ortholattices*, Ph.D. thesis, TU Darmstadt, January 2003, <http://elib.tu-darmstadt.de/diss/000303/diss.pdf>.

- [13] F. Wehrung, A uniform refinement property for congruence lattices, Proc. Amer. Math. Soc. **127** (1999), 363–370.

(C. Herrmann) TUD FB4, SCHLOSSGARTENSTR. 7, 64289 DARMSTADT, GERMANY

E-mail address: herrmann@mathematik.tu-darmstadt.de