

# PERSPECTIVITY IN COMPLEMENTED MODULAR LATTICES AND REGULAR RINGS

CHRISTIAN HERRMANN

ABSTRACT. Based on an analogue for systems of partial isomorphisms between lower sections in a complemented modular lattice we construct a series of terms (including inner inverse as basic operation and providing descending chains) such that principal right ideals  $aR \cong bR$  in a (von Neumann) regular ring  $R$  are perspective if the series becomes stationary. In particular, this applies if  $aR \cap bR$  is of finite height in  $L(R)$ . This is used to derive, for existence-varieties  $\mathcal{V}$  of regular rings, equivalence of unit-regularity and direct finiteness, both conceived as a property shared by all members of  $\mathcal{V}$ .

## 1. INTRODUCTION

(Von Neumann) regular rings  $R$  and complemented modular lattices are closely connected fields since the work of von Neumann cf. [26] - with  $R$  one associates its lattice  $L(R)$  of principal right ideals. *Unit-quasi-inverses*  $u$  of elements  $a$  (i.e. unites  $u$  such that  $aua = a$ ) have been introduced by Ehrlich [6, 7], a ring being *unit-regular* if each element admits some unit-quasi-inverse (such rings are, in particular *directly finite*:  $ab = 1$  implies  $ba = 1$ ). Ehrlich also showed that a regular ring  $R$  is unit-regular if and only if for all idempotents  $e, f$  one has  $eR \cong fR$  implying  $(1 - e)R \cong (1 - f)R$ . Handelman [13] added further equivalent conditions, one of them being that  $eR \cong fR$  implies  $eR$  perspective to  $fR$  in  $L(R)$ . Perspectivity, regularity, and unit-regularity of elements in general rings have been intensively studied, see e.g. [8, 21, 22, 24].

The purpose of the present note is to give a sufficient condition on  $aR \cong bR$  in a regular ring  $R$  granting that  $aR$  is perspective to  $bR$  (which holds if  $a$  has a unit-quasi-inverse) and to show that this applies if  $aR \cap bR$  is of finite height in  $L(R)$ .

---

1991 *Mathematics Subject Classification.* 06C20, 16E50.

*Key words and phrases.* Complemented modular lattice, von Neumann regular ring, perspectivity, unit-regularity, existence-varieties.

Here, establishing perspectivity relies on calculations in  $L(R)$ , for convenience done in abstract complemented modular lattices endowed with a system of isomorphisms between lower sections requiring properties present in the case of isomorphisms induced by isomorphisms between principal right ideals. The principal result is a reduction process associating  $e_{n+1} \leq e_n$  and  $f_{n+1} \leq f_n$  with given  $e_n, f_n$  such that  $e_n$  is perspective to  $f_n$  (and so  $e_0$  perspective to  $f_0$ ) if  $e_{n+1} \cap f_{n+1} = e_n \cap f_n$  or if  $e_{n+1}$  is perspective to  $f_{n+1}$ . In  $\aleph_0$ -complete complemented modular lattices  $e_0$  is perspective to  $f_0$  if the meet of the  $e_n$  is perspective to the meet of the  $f_n$ .

If one considers regular rings endowed with an operation of inner inversion, termination of this reduction process after  $n$  steps can be captured by an identity. This is applied to study unit-regularity in the context of *existence varieties*  $\mathcal{V}$  of regular rings, that is, classes closed under homomorphic images, direct products, and regular subrings. It is shown that for such classes unit-regularity is equivalent to direct finiteness, both considered as a property required for all members. (Compare this to the result of Baccella and Spinosa [3] that a semiartinian regular ring is unit-regular if and only if all its homomorphic images are directly finite.) Another property shown equivalent to unit-regularity is that  $\mathcal{V}$  does not contain nonartinian subdirectly irreducibles, equivalently, if it is generated by artinians of bounded finite length. Having  $\mathcal{V}$  generated by artinians is not sufficient for unit-regularity in view of the result, established by Goodearl, Menal, and Moncasi [10, Thm. 2.5], that free regular rings are residually artinian (and, according to Herrmann and Semenova [15, Cor. 14], even residually finite). These results further developed work of Tyukavkin [27] obtaining the ring of row and column finite matrices as well as certain regular rings  $R$  of endomorphisms of vector spaces as homomorphic images of (regular) subrings of products of finite-dimensional matrix rings over (skew-)fields. In particular, this approach was basic for the study of existence varieties of regular rings in Herrmann and Semenova [15] and of varieties of  $*$ -regular rings which are generated by their artinian members, see Micol [23], Herrmann and Semenova [16], and [18].

Thanks are due to the referees for valuable hints to related literature and results as well as to necessary clarifications and a possible extension.

## 2. COMPLEMENTED MODULAR LATTICES

**2.1. Preliminaries.** We refer to Birkhoff [5] and von Neumann [26]. A *lattice*  $L$  is a set endowed with a partial order  $\leq$  such that any two

elements  $a, b$  have infimum and supremum written as *meet*  $a \cap b$  and *join*  $a + b$ , respectively. We also write  $ab = a \cap b$  and apply the usual preference rules. All lattices to be considered will have smallest element 0 and greatest element 1.

For  $u \leq v$  in  $L$ , the *interval*  $[u, v] = \{x \mid u \leq x \leq v\}$  is again a lattice with the inherited partial order and operations. A lattice  $L$  is *modular* if

$$b \leq a \Rightarrow a(b + c) = b + ac.$$

Then the maps  $x \mapsto x + b$  and  $y \mapsto ay$  are mutually inverse isomorphisms between  $[ab, a]$  and  $[b, b + a]$ . An element  $u$  of a modular lattice  $L$  is *neutral* if  $(u + x)(u + y) = u + xy$  for all  $x, y \in L$ ; the set of neutral elements is a sublattice of  $L$ . An element  $a$  of a modular lattice  $L$  is of *height*  $d$  if some (equivalently: each) maximal chain in  $[0, a]$  has *length*  $d$ , that is  $d + 1$ -elements. For the following see [5, Ch. III Thm. 15].

**Fact 1.** *In a modular lattice, the direct product  $[ab, a] \times [ab, b]$  embeds into  $[ab, a + b]$  via  $(x, y) \mapsto x + y = (x + b)(y + a)$ .*

If  $ab = 0$  then we write  $a + b = a \oplus b$ . If  $a \oplus b = 1$  then  $b$  is a *complement* of  $a$ . A lattice  $L$  is *complemented* if each element admits some complement. If  $L$  is, in addition, modular then we speak of a CML. In a CML each interval  $[u, v]$  is again a CML (within  $[u, v]$ , a complement of  $x$  is given by  $yv + u = (y + u)v$  where  $x \oplus y = 1$  in  $L$ ).

In a CML, elements  $a, b$  are *perspective*, written as  $a \sim b$ , if they have a common complement; equivalently,  $a \sim_c b$  for some  $c$ , the latter meaning that  $a + b = a + c = b + c$  and  $ab = ac = bc$ . We write  $a \approx_c b$  if  $a \sim_c b$  and  $ab = 0$ ; also  $a \approx b$  if  $a \approx_c b$  for some  $c$ . Applying Fact 1 one obtains the following.

**Fact 2.** *In a modular lattice, if  $a_i \sim_{c_i} b_i, i = 1, 2$  and  $a_1 b_1 a_2 b_2 \geq (a_1 + b_1)(a_2 + b_2)$  then  $a_1 + b_1 \sim_{c_1 + c_2} a_2 + b_2$ .*

**Fact 3.** *In a modular lattice one has  $a \sim b$  if and only if  $x \sim y$  for some (equivalently: all)  $x, y$  such that  $a = x \oplus ab$  and  $b = y \oplus ab$ . Moreover, one has  $a \oplus y = a + b = b \oplus x$  for such  $x, y$ .*

*Proof.* Observe that for such  $x, y$  one has  $ay = aby = 0$  whence by modularity  $a(x + y) = x$  and, similarly,  $bx = 0$  and  $b(x + y) = y$ . By modularity it follows that  $ab(x + y) = 0$ . Now one has  $ab + x + y = a + b$  so that the map  $z \mapsto z + ab$  is an isomorphism of  $[0, x + y]$  onto  $[ab, a + b]$ . Moreover,  $a + y = a + ab + y = a + b$ . Thus  $a \oplus y = a + b$  and, similarly,  $b \oplus x = a + b$ .  $\square$




FIGURE 1. Lemma 4

## 2.2. Two lemmas on modular lattices.

**Lemma 4.** *In a modular lattice, if  $z = x \oplus y$ ,  $w = u \oplus v$ , and  $zw = xu$  then  $yv = 0$ . If, in addition,  $x \sim u$  and  $y \sim v$  then also  $z \sim w$ .*

*Proof.* Clearly,  $yv \leq yzw \leq yx = 0$ . Moreover, by modularity one has  $(x + w)z = x + wz = x$  whence  $(x + w)y = (x + w)zy = xy = 0$  and, similarly,  $(u + z)v = 0$ . Now, by Fact 1  $(x + w)(u + z) = x + u$  it follows  $(y + v)(x + u) = (y + v)(x + w)(u + z) = [y(x + w) + v](u + z) = v(u + z) = 0$  by modularity. In particular, this implies  $xu(y + v) = 0$  establishing the isomorphism  $r \mapsto r + xu$  of  $[0, y + v]$  onto  $[xu, y + xu + v]$ . Thus, if  $y \sim v$  then one has also  $y + xu \sim v + xu$ . Assuming that, in addition,  $x \sim u$  one derives  $z \sim w$  by Fact 2 since  $(x + u)(y + xu + v) = (x + u)(y + v) + xu = xu$ .  $\square$

**Lemma 5.** *In modular lattice,  $L$ ,  $a \sim_c b$  and  $d = (a + d)(b + d)$  jointly imply  $a + d \sim b + d$ .*

*Proof.* Assume  $a \sim_c b$  in  $L$  and let  $S$  denote the sublattice  $S$  generated by  $a, b, c, d$ . Let  $D_2$  and  $M_3$  denote the 2-element lattice and the height 2 lattice with 3 atoms, respectively. Obviously,  $S$  is also generated by  $c$ , and the two chains  $a \leq a + d$  and  $b \leq b + d$ . Thus, by [14]  $S$  is a sub direct product of lattices  $D_2$ ,  $M_3$ , and lattices  $M$  where (the images of)  $a, a + d, b, b + d$  generate a boolean sublattice with  $(a * b)(a + d)(b + d) = 0$  and  $a + d + b = 1$  and where (the image of)  $c$  is a common complement of the “atoms”  $d = (a + d)(b + d)$ ,  $a$ , and  $b$ . Since also  $a \sim_c b$ ,  $M$  is easily seen to be trivial in both cases..




FIGURE 2. Lemma 5

Thus,  $S$  is a subdirect product of factors  $D_2$  and  $M_3$ , only. Due to the given relations, in any of these factors the images of  $d$  and  $a + b$  take value 0 or 1, only; this means that  $a + b$  and  $d$  are neutral elements of  $S$ . It follows, that  $u = d(a + b)$  is neutral, too, whence  $a \sim_c b$  implies  $a + u \sim_{c+u} b + u$  and this in turn  $a + d \sim_{c+d} b + d$  via the isomorphism of  $[u, a + b]$  onto  $[d, 1]$ .  $\square$

**2.3. Partial isomorphisms and perspectivity.** Motivated by the case where  $L$  is the lattice of principal right ideals of a regular ring, for any CML  $L$  we consider lattice isomorphisms  $\alpha : [0, e] \rightarrow [0, f]$ , shortly written as  $\alpha : e \rightarrow f$ . For  $g = e \cap f$  one has both  $\alpha^{-1}(g)$  and  $\alpha(g)$  defined. If  $e' \leq e$  let  $\alpha|_{e'}$  denote the restriction of  $\alpha$  to  $[0, e']$ , that is  $\alpha|_{e'} : e' \rightarrow f'$ ,  $f' = \alpha(e')$ .

**Observation 6.**  $\alpha^{-1}(g) \cap \alpha(g) \leq g$  and equality holds if and only if  $\alpha(g) = g$ .

*Proof.* Clearly,  $\alpha^{-1}(g) \cap \alpha(g) \leq \alpha^{-1}(f) \cap \alpha(e) = e \cap f = g$ . Also, if  $\alpha(g) = g$  then  $\alpha^{-1}(g) = g$  whence  $\alpha^{-1}(g) \cap \alpha(g) = g$ . Conversely, if the latter holds then  $g \leq \alpha(g)$  and  $\alpha(g) \leq \alpha(\alpha^{-1}(g)) = g$  and it follows  $g = \alpha(g)$ .  $\square$

We introduce a reduction process which yields perspectivity provided that it stops after finitely many steps. With  $g = e \cap f$  one has  $\alpha^\# : \alpha^{-1}(g) \rightarrow \alpha(g)$  defined by  $\alpha^\#(x) = \alpha^2(x)$ .

We consider *admissible* systems  $A$  of isomorphisms  $\alpha : e \rightarrow f$  requiring the following axioms:

- (A1) If  $\alpha : e \rightarrow f$  is in  $A$  and  $e' \leq e$  then  $\alpha|_{e'}$  is in  $A$ .

(A2) If  $\alpha : e \rightarrow f$  is in  $A$  and  $e \cap f = 0$  then  $e \approx f$ .

(A3) If  $\alpha : e \rightarrow f$  is in  $A$  and  $g = e \cap f$  then  $\alpha^\#$  is in  $A$ .

**Lemma 7.** *Consider  $\alpha : e \rightarrow f$  in admissible  $A$ .*

(i) *If  $\alpha(e \cap f) = e \cap f$  then  $e \sim f$ .*

(ii) *If  $\alpha^{-1}(e \cap f) \sim \alpha(e \cap f)$  then  $e \sim f$ .*

*Proof.* Let  $g = e \cap f$ . In (i) choose  $y$  so that  $y \oplus g = e$  and put  $v = \alpha(y)$ . Then  $f = \alpha(e) = \alpha(y \oplus g) = \alpha(y) \oplus \alpha(g) = v \oplus g$ . With  $x = u = g$  in Lemma 4 it follows  $yv = 0$  whence  $y \sim v$  by axioms (A1) and (A2); moreover,  $e \sim f$  by Lemma 4, again.

(ii): Put  $x = g + \alpha^{-1}(g)$  and  $u = g + \alpha(g)$ , observe that  $u = \alpha(x)$ . Observe that  $g \leq x \leq e$  and  $g \leq u \leq f$  whence  $x \cap u = g$ . Thus, by hypothesis and Lemma 5 one has  $x \sim u$ . Choose  $y$  such that  $y \oplus x = e$  and put  $v := \alpha(y)$ . Then  $f = \alpha(x \oplus y) = \alpha(x) \oplus v$  and we conclude  $y \cap v = 0$  by Lemma 4 whence  $y \sim v$  by axioms (A1) and (A2); finally,  $e \sim f$  by Lemma 4.  $\square$

Given  $\alpha : e \rightarrow f$  in admissible  $A$  define, by induction,  $\alpha_0 = \alpha$ ,  $e_0 = e$ ,  $f_0 = f$  and  $\alpha_{n+1} = \alpha_n^\# : e_{n+1} \rightarrow f_{n+1}$ . Put  $g_n = e_n \cap f_n$  and observe that  $e_{n+1} \leq e_n$  and  $f_{n+1} \leq f_n$  whence also  $g_{n+1} \leq g_n$ .

**Theorem 8.** *Given a CML,  $L$ , an admissible system  $A$  of partial isomorphisms in  $L$ , and  $\alpha : e \rightarrow f$  in  $A$ , one has  $e \sim f$  provided that  $e_m \sim f_m$  for some  $m$ . In particular, this applies if  $\alpha_m(g_m) = g_m$  respectively  $g_{m+1} = g_m$  for some  $m$  or if  $e \cap f$  is of finite height in  $L$ .*

*Proof.* If  $e_m \sim f_m$  (which by (i) of Lemma 7 is the case if  $\alpha_m(g_m) = g_m$ ) then  $e_{m-1} \sim f_{m-1}$  by (ii) of Lemma 7 and it follows  $e \sim f$  by induction. Now, assume  $e \cap f$  of finite height in  $L$ . By Observation 6,  $g_{m+1} < g_m$  unless  $\alpha_m(g_m) = g_m$ ; thus, the latter has to occur for some  $m$ .  $\square$

We now give for the case  $g_{m+1} = g_m$  a proof with a single application of Lemma 5, only. A sequence  $a_0, \dots, a_m$  in a modular lattice is *independent*, if for all  $n < m$ ,  $a_n \sum_{n < k \leq m} a_k = 0$ ; equivalently if  $(\sum_{n \in I} a_n)(\sum_{n \in J} a_n) = 0$  for all  $I, J \subseteq \{0, \dots, m\}$  such that  $I \cap J = \emptyset$ . Induction using Fact 2 yields the following.

**Fact 9.** *In a modular lattice, if  $a_0 + b_0, a_1 + b_1, \dots, a_m + b_m$  is independent and  $a_n \approx b_n$  for all  $n \leq m$  then  $\sum_{n \leq m} a_n \approx \sum_{n \leq m} b_n$ .*

**Lemma 10.** *Consider  $\alpha_n : e_n \rightarrow f_n$  and  $g_n$ ,  $n \leq m + 1$  as above and, for  $n \leq m$ ,  $x_n$  such that  $e_n = x_n \oplus (e_{n+1} + g_n)$  and  $y_n = \alpha_n(x_n)$ . Then the following hold.*

(i)  *$f_n = y_n \oplus (g_n + f_{n+1})$  and  $(e_k + f_k)(x_n + y_n) = 0$  for all  $k > n$ .*

(ii)  *$x_n \approx y_n$  for all  $n \leq m$ .*

- (iii)  $x_0, y_0, \dots, x_m, y_m$  is an independent sequence.
- (iv)  $x := \sum_{k \leq m} x_k \approx y := \sum_{k \leq m} y_k$ .
- (v) If  $g_m = g_{m+1}$  then  $x + g_0 = e_0$ ,  $y + g_0 = f_0$ , and  $e_0 \sim f_0$ .

*Proof.* Recall that  $e_{n+1} = \alpha_n^{-1}(g_n) \leq e_n$  and  $f_{n+1} = \alpha_n(g_n) = \alpha_n^2(e_{n+1}) \leq f_n$  and observe that  $e_n = x_n \oplus (\alpha_n^{-1}(g_n) + g_n)$ . It follows that  $f_n = \alpha_n(e_n) = y_n \oplus (g_n + \alpha_n(g_n)) = y_n \oplus (g_n + f_{n+1})$  for  $n \leq m$ . Observe that  $x_n y_n \leq e_n f_n = g_n$  whence  $x_n y_n \leq x_n g_n = 0$ . Thus  $x_n \approx y_n$  by axiom (A2).

Moreover, modularity yields  $(e_{n+1} + f_{n+1})(x_n + y_n) \leq (e_{n+1} + f_n)(x_n + y_n) = (e_{n+1} + f_n)x_n + y_n = (e_{n+1} + f_n)e_n x_n + y_n = (e_{n+1} + f_n e_n)x_n + y_n = (e_{n+1} + g_n)x_n + y_n = 0 + y_n = y_n$  and so  $(e_{n+1} + f_{n+1})(x_n + y_n) = (e_{n+1} + f_{n+1})y_n = (e_{n+1} + f_{n+1})f_n y_n = (e_{n+1} f_n + f_{n+1})y_n \leq (g_n + f_{n+1})y_n = 0$ . It follows for all  $n < m$

$$\left( \sum_{n < k \leq m} x_k + y_k \right) (x_n + y_n) \leq (e_{n+1} + f_{n+1})(x_n + y_n) = 0,$$

proving (iii); (iv) follows by Fact 9. Dealing with (v), observe that  $e_{m+1} = g_m$  whence  $g_0 + x_m \geq e_m$ . Now, backward induction yields  $g_0 + \sum_{m \geq k \geq n} x_k \geq e_n$  for all  $n \leq m$ , whence  $g_0 + x = e_0$ . Similarly, one has  $g_0 + y = f_0$  and it follows  $e_0 \sim f_0$  by Lemma 2.  $\square$

**2.4.  $\aleph_0$ -complete complemented modular lattices.** A CML is  $\aleph_0$ -complete if supremum  $\sum_n a_n$  and infimum  $\prod_n a_n$  exist for all families  $(a_n \mid n < \omega)$ . According to Amemiya and Halperin [1, Thm.9.5] any such is also  $\aleph_0$ -continuous, i.e.  $b \sum_n a_n = \sum_n b a_n$  (resp.  $b + \prod_n a_n = \prod_n (b + a_n)$ ) if  $(a_n \mid n < \omega)$  is upward (resp. downward) directed. A sequence  $(a_n \mid n < \omega)$  is *independent* if each of its finite subsequences is independent.

**Fact 11.** *In an  $\aleph_0$ -complete CML, if  $a_0 + b_0, a_1 + b_1, \dots$  is independent and  $a_n \approx b_n$  for all  $n < \omega$  then  $\sum_n a_n \approx \sum_n b_n$ .*

*Proof.* Suppose  $a_n \approx_{c_n} b_n$  for  $n < \omega$  and write  $x_n^+ = \sum_{m \leq n} x_m$  and  $x_\omega = \sum_{n < \omega} x_n = \sum_{n < \omega} x_n^+$  for any sequence  $x_0, x_1, \dots$ . By Fact 9 one has  $a_n^+ \approx_{c_n^+} b_n^+$  for all  $n$ . Since sequences  $x_0^+, x_1^+, \dots$  are upward directed,  $\aleph_0$ -continuity yields

$$\begin{aligned} a_\omega b_\omega &= \left( \sum_{n < \omega} a_n^+ \right) b_\omega = \sum_{n < \omega} a_n^+ b_\omega = \sum_{n < \omega} a_n^+ \sum_{m < \omega} b_m^+ = \\ &= \sum_{n < \omega} \sum_{m < \omega} a_n^+ b_m^+ \leq \sum_{n, m < \omega} a_{\max(n, m)}^+ b_{\max(n, m)}^+ = 0. \end{aligned}$$

By symmetry one obtains  $a_\omega c_\omega = b_\omega c_\omega = 0$  while  $a_\omega + b_\omega = a_\omega + c_\omega = b_\omega + c_\omega$  is obvious.  $\square$

**Theorem 12.** *Assume that  $\alpha : e \rightarrow f$  is member of an admissible system  $A$  of partial isomorphisms of an  $\aleph_0$ -complete CML,  $L$ , and that  $\prod_{n < \omega} e_n \sim \prod_{n < \omega} f_n$  for  $e_n, f_n$  defined as in Subsection 2.3. Then it follows that  $e \sim f$  in  $L$ .*

*Proof.* Given  $\alpha : e \rightarrow f$  in  $A$  define  $\alpha_n : e_n \rightarrow f_n$  as in Subsection 2.3 and  $g_n = e_n f_n$ . Put  $e_\infty = \prod_n e_n$ ,  $f_\infty = \prod_n f_n$ , and  $g_\infty = \prod_n g_n = e_\infty f_\infty$  and recall that  $e_{n+1} = \alpha_n^{-1}(g_n) \leq e_n$  and  $f_{n+1} = \alpha_n(g_n) = \alpha_n^2(e_{n+1}) \leq f_n$ .

Choose  $x_n$  such that  $e_n = x_n \oplus (e_{n+1} + g_n)$  and  $y_n = \alpha_n(x_n)$ . By Lemma 10 one has  $f_n = \alpha_n(e_n) = y_n \oplus (g_n + f_{n+1})$  and the sequence  $x_0, y_0, x_1, y_1, \dots$  is independent; thus, Fact 11 yields  $x_\omega \approx y_\omega$  where  $x_\omega = \sum_n x_n$  and  $y_\omega = \sum_n y_n$ . On the other hand, again by Lemma 10 and modularity,

$$\left( e_\infty + f_\infty + \sum_{n < k \leq m} x_k + y_k \right) (x_n + y_n) = 0 \text{ for all } m > n$$

that is  $e_\infty + f_\infty, x_0 + y_0, x_1 + y_1, \dots$  is an independent sequence, too. By hypothesis  $e_\infty \sim f_\infty$  and in view of  $(x_\omega + y_\omega)(e_\infty + f_\infty) = 0$  Fact 2 applies to yield  $x_\omega + e_\infty \sim y_\omega + f_\infty$ .

Now  $g + e_{n+1} + x_\omega \geq e_{n+1} + g_n + x_n = e_n$  and by induction it follows  $g + e_n + x_\omega = e$  for all  $n$ . Thus, by  $\aleph_0$ -continuity one has  $g + x_\omega + e_\infty = \prod_n (g + x_\omega + e_n) = e$ . Similarly, one obtains  $g + y_\omega + f_\infty = f$ . Finally,  $e \sim f$  follows by Lemma 5.  $\square$

### 3. REGULAR RINGS

**3.1. Preliminaries.** A ring  $R$  (associative and with unit) is (von Neumann) *regular* if for each  $a \in R$  there is a *quasi-inverse* or *inner inverse*  $x \in R$  such that  $axa = a$ ; equivalently, every right (left) principal ideal is generated by an idempotent, see Goodearl [9] and Wehrung [28]. For a regular ring,  $R$ , the principal right ideals form a complemented sublattice  $L(R)$  of the lattice of all right ideals; in particular,  $L(R)$  is modular. For artinian  $R$ , the height of  $L(R)$  is the *length* of  $R$ .

An element  $a$  of  $R$  is *unit-regular* if there is a unit  $u \in R$ , a *unit-quasi-inverse*, such that  $aua = a$ .  $R$  is *unit-regular* if all its elements are unit-regular. Any such ring is *directly finite* (that is  $ab = 1$  implies  $ba = 1$ ), the converse not being true for regular rings, in general.



If  $e$  is an idempotent in a regular ring  $R$ , then the *corner*  $eRe$  is a regular ring with unit  $e$ , a homomorphic image of the regular subring  $eRe + (1 - e)R(1 - e)$  of  $R$ .

Idempotents  $e, f$  are Murray von Neumann *equivalent* if  $e = yx$  and  $f = xy$  for some  $x, y$ . For the following see e.g. Handelman [13] and Goodearl [9].

- (1) For any  $a$  there is a *generalized* or *reflexive inverse*  $b$  such that  $aba = a$  and  $bab = b$ , e.g.  $b = xax$  where  $axa = a$ . Then  $ab$  and  $ba$  are idempotents.
- (2)  $e, f$  are idempotents and equivalent if and only if  $e = ba$  and  $f = ab$  for some  $a, b$  as in (1). Moreover, in this case  $\omega_{a,b}(r) = ar$  defines an isomorphism  $\omega_{a,b} : bR = eR \rightarrow aR = fR$  of right  $R$ -modules with inverse  $\omega_{b,a}$ . It follows that  $fae = a$  and  $ebf = b$ .
- (3) For idempotents  $e, f$ , every  $R$ -module isomorphism  $\omega : eR \rightarrow fR$  is as in (2) where  $\omega(e) = a$  and  $\omega^{-1}(f) = b$ .
- (4) If  $eR \sim fR$  in  $L(R)$  then  $eR \cong fR$  as right  $R$ -modules. If  $eR \cap fR = 0$  then the converse holds, too.
- (5) If  $c$  is another generalized inverse of  $a$  then  $bR \sim cR$  (being complements of  $\{x \in R \mid ax = 0\}$ ) and  $x \mapsto cax$  is an isomorphism of  $bR$  onto  $cR$ .

To prove (3), put  $a = \omega(e)$  and  $b = \omega^{-1}(f)$ . Then one has  $\omega(r) = \omega(er) = \omega(e)r = ar$  for  $r \in eR$  and, similarly,  $\omega^{-1}(s) = bs$  for  $s \in fR$ . Thus,  $aba = \omega(ba) = \omega(\omega^{-1}(a)) = a$  and, similarly,  $bab = b$ .

A regular ring  $R$  is *perspective* if isomorphic direct summands of  $R_R$  are perspective in  $L(R)$ ; equivalently,  $aR$  is perspective to  $bR$  for all  $aR \cong bR$  - for a more general result see Mary [22, Thm.3.1].

**Theorem 13.** Handelman [13] *A regular ring is unit-regular if and only if it is perspective.*

An element  $a$  of  $R$  is *strongly  $\pi$ -regular* if there is  $n$  such that  $a^n \in a^{n+1}R \cap Ra^{n+1}$ .  $R$  is *strongly  $\pi$ -regular* if so are all its elements.

**Theorem 14.** Goodearl and Menal [12, Thm. 5.8]. *Strongly  $\pi$ -regular regular rings are unit-regular.*

In general rings, a strongly  $\pi$ -regular element is unit-regular provided all its powers are regular. For a detailed discussion and proofs see Khurana [20]

**3.2. Perspectivity.** The following “local version” of Handelman’s Theorem should be well known.

**Lemma 15.** *Given a generalized inverse  $b$  of  $a$  in a regular ring  $R$  one has idempotents  $e = ba$ ,  $f = ab$ , and  $g$  such that  $gR = eR \cap fR$ . Now, the following are equivalent.*

- (i)  $fR$  and  $eR$  are perspective in  $L(R)$ .
- (ii) For some (all) idempotents  $e', f'$  such that  $e'R \oplus gR = eR$  and  $f'R \oplus gR = fR$  one has  $e'R \cong f'R$ .
- (iii) For some (all) idempotents  $e', f'$  such that  $e'R \oplus gR = eR$  and  $f'R \oplus gR = fR$  there is a unit  $u$  of  $R$  such that  $aua = a$  and  $e'R \cong f'R$  via  $ue'$ .

*Proof.* In view of (2) in the preceding subsection,  $e = ba$  and  $f = ab$  are idempotents. Consider  $e', f'$  as in (ii) and (iii) and observe that such exist since  $L(R)$  is complemented.

By Fact 3 we have  $e'R \cap f'R = 0$  and, moreover,  $eR \sim fR$  if and only if  $e'R \sim f'R$ . By (4), the latter is equivalent to  $e'R \cong f'R$ . This proves that (i) is equivalent to (ii).

Now, assume (ii), in particular  $f'R \cong e'R$  via some isomorphism  $\omega'$ . Choose an idempotent  $h$  such that  $eR + fR = hR$ . Then

$$eR \oplus f'R \oplus (1-h)R = R = fR \oplus e'R \oplus (1-h)R,$$

again by Fact 3. In view of (2) define

$$\omega(r + s + t) = \omega_{b,a}(r) + \omega'(s) + t \text{ for } r \in fR, s \in e'R \text{ and } t \in (1-h)R$$

to obtain an automorphism of the right  $R$ -module  $R = 1R$ . By (3) there are  $u, v$  in  $R$  such that  $\omega = \omega_{u,v}$ ; in particular,  $u$  is a unit and  $v = u^{-1}$ . Moreover,  $ur = \omega_{b,a}(r) = br$  for  $r \in fR$ , in particular  $ua = ba = e$ . Thus  $aua = ae = a$ , proving that (ii) implies (iii).

Finally, assume (iii). Then  $u^{-1}f' = e'$  whence  $x \mapsto ux$  is an  $R$ -module isomorphism of  $e'R$  onto  $f'R$  with inverse  $y \mapsto u^{-1}y$ . Thus, (ii) and (iii) are equivalent, too.  $\square$

For  $A, B \in L(R)$  and right module isomorphism  $\omega : A \rightarrow B$  one has the induced lattice isomorphism  $\omega_L : [0, A] \rightarrow [0, B]$ . Let  $A(R)$  denote the set of all these.

**Lemma 16.**  *$A(R)$  is an admissible system of partial isomorphisms of  $L(R)$ .*

*Proof.* Consider  $\omega : A \rightarrow B$  in  $A(R)$  and observe that  $\omega_L(X) = \omega(X)$  for all  $X \leq A$ . Thus, if  $A' \leq A$  in  $L(R)$  then  $(\omega|_{A'})_L : [0, A'] \rightarrow [0, B']$  in  $A(R)$  with  $B' = \omega(A') \leq B$ , proving axiom (A1). Similarly, for  $C = A \cap B$ ,  $A' = \omega^{-1}(C)$ , and  $B' = \omega(C)$  one has  $\omega_L^\# = (\omega|_C \circ \omega|_{A'})_L$  in  $A(R)$ , proving axiom (A3). Finally, (A2) follows from (4).  $\square$

As observed by the referee, the following can be obtained in purely ring theoretic terms: it is an immediate consequence of Lemma 15 and [9, Proposition 4.13].

**Corollary 17.** *For  $A, B$  in the lattice  $L(R)$  of principal right ideals of the regular ring  $R$ , if  $A \cap B$  is of finite height in  $L(R)$  then  $A, B$  are perspective in  $L(R)$  if and only if they are isomorphic as  $R$ -modules.*

*Proof.* Assume  $A \cap B$  of finite height. If  $A \cong B$  then there is a lattice isomorphism  $\alpha : [0, A] \rightarrow [0, B]$  in  $A(R)$  and in view of Lemma 16 and Theorem 8 it follows that  $A \sim B$ . The converse follows from (4).  $\square$

**Corollary 18.** *An element  $a$  in a regular ring  $R$  is unit-regular provided that there is a reflexive inverse  $b$  of  $a$  such that  $bR \cap aR$  is of finite height in  $L(R)$ .*

**3.3. Regular rings with operation of quasi-inversion.** A regular ring may be considered as an algebraic structure also endowed with an operation  $a \mapsto a'$  of quasi-inversion. The class  $\mathcal{R}$  of all these structures is then defined by the identities for rings with unit together with  $xx'x = x$ . As observed above, the term  $x^+ = x'xx'$  then yields a generalized inverse  $a^+$  of  $a$  and  $\gamma(x) = xx^+$  yields idempotents  $\gamma(a)$  such that  $\gamma(a)R = aR$ . Though, in this setting, regular subalgebras (these algebras endowed with an operation of inner inversion) of unit-regular rings may fail to be unit-regular as shown by Bergman [4]. As remarked by the referee, unit-regularity can be equationally defined adding the identities  $x'(x')' = 1$  and  $(x')'x' = 1$ . See Ara, Goodearl, Nielsen, Pardo, and Perera [2] for a comprehensive introduction.

For the following see Wehrung [28, Lemma 8-3.12].

**Lemma 19.** *There are binary terms  $x \vee y$ ,  $x \wedge y$ , and  $x \ominus y$  in the language of  $\mathcal{R}$  such that, for all  $R \in \mathcal{R}$  and  $a, b \in R$ ,  $a \vee b$ ,  $a \wedge b$ , and  $a \ominus b$  are idempotent,  $(a \vee b)R = aR + bR$ ,  $(a \wedge b)R = aR \cap bR$ , and  $(a \ominus (a \wedge b))R \oplus (a \wedge b)R = aR$ .*

**Theorem 20.** *For each natural number  $n$  there are binary terms  $t_n(x, y)$ ,  $u_n(x, y)$ , and  $p_n(x, y)$  in the language of  $\mathcal{R}$  such that the following hold for all  $R \in \mathcal{R}$  and mutually reflexive inverses  $a, b \in R$ :  $t_n(a, b)$  is idempotent; moreover, if  $t_{n+1}(a, b)t_n(a, b) = t_n(a, b)$  then*

- (i)  $bR$  and  $aR$  are perspective in  $L(R)$ :  $bR \sim_{p_n(a, b)} aR$ .
- (ii)  $u_n(a, b)$  is a unit such that  $au_n(a, b)a = a$ .

*If  $R$  is of length at most  $n + 2$  then  $t_{n+1}(a, b)t_n(a, b) = t_n(a, b)$  for all mutually reflexive inverses  $a, b$  in  $R$ .*

*Proof.* With idempotents  $e_0 = ba$  and  $f_0 = ab$  one has the isomorphism  $\omega_{ab} : e_0R \rightarrow f_0$  given by  $x \mapsto ax$  inducing the isomorphism  $\alpha = \alpha_0 :$

$[0, eR] \rightarrow [0, fR]$  given by  $\alpha(xR) = axR$  with inverse  $\alpha^{-1}(xR) = bxR$ . Recalling the construction in Subsection 2.3 put  $g_0 = e_0 \wedge f_0$  and, recursively,

$$g_n = e_n \wedge f_n, e_{n+1} = \gamma(b^{2^n} g_n), f_{n+1} = \gamma(a^{2^n} g_n)$$

to obtain  $\alpha_{n+1} : [0, e_{n+1}R] \rightarrow [0, f_{n+1}R]$  given by  $\alpha_{n+1}(xR) = \alpha_n^2(xR) = a^{2^n} xR$ . Accordingly, put  $t_0(x, y) = yx \wedge xxy$ , and, inductively,

$$t_{n+1}(x, y) = y^{2^n} t_n(x, y) \wedge x^{2^n} t_n(x, y).$$

Thus, for  $a, b$  as above one has  $t_n(a, b)R = g_nR = e_nR \cap f_nR$  whence

$$t_{n+1}(a, b)t_n(a, b) = t_n(a, b) \Leftrightarrow g_{n+1}R = g_nR.$$

Thus, supposing  $t_{n+1}(a, b)t_n(a, b) = t_n(a, b)$ ,  $bR$  and  $aR$  are perspective in  $L(R)$  by Theorem 8 and Lemma 15 applies to provide the existence of a unit  $u$  in  $R$  such that  $aua = a$  and idempotent  $p \in R$  such that  $bR \sim_{pR} aR$ . To prove the existence of terms  $u_n(x, y)$  and  $p_n(x, y)$ , as required, it suffices to observe that all this applies, in particular, to  $R$  being the free algebra in  $\mathcal{R}$  with generators  $a, b$  and relations  $aba = a$ ,  $bab = b$ , and  $t_{n+1}(a, b)t_n(a, b) = t_n(a, b)$ .

Now, assume that  $g_k \neq g_{k+1}$  for all  $k \leq m$ . Then one obtains a chain  $e_0R + f_0R > e_0R > g_0R > \dots > g_{m+1}R$  of length  $m+3$  in  $L(R)$ . Thus, if  $r$  is of length at most  $n+2$  then  $g_m = g_{m+1}$  for some  $m \leq n+2$  and it follows  $g_k = g_m$  for all  $k \geq m$ , in particular  $g_n = g_m = g_{n+1}$ .  $\square$

**Example 21.** (i) *There are  $a, b, c$  in some unit-regular ring  $R$  such that  $a, b$  and  $a, c$  are pairs of reflexive inverses,  $aR, bR$ , and  $cR$  pairwise perspective,  $t_{n+1}(a, b)t_n(a, b) \neq t_n(a, b)$  for all  $n$ , and  $t_0(a, c) = 0$ .*

(ii) *There are a regular ring  $R$  and reflexive inverses  $a, b$  in  $R$  such that  $t_0(a, b) = 0$  but both  $a$  and  $b$  are not strongly  $\pi$ -regular,*

*Proof.* Considering (i) let  $V$  a vector space of dimension  $n+3$ . We show by induction that  $\mathbf{End}(V)$  contains some  $a, b$  with associated  $g_n > g_{n+1}$ . More precisely, we show that for any subspaces  $V_1 \neq V_2$  of codimension 1 there is such  $a$  with generalized inverse  $a^+$  and restricting to an isomorphism  $V_1 \rightarrow V_2$  and such that  $V_1 = \mathbf{im} a^+$  and  $V_2 = \mathbf{im} a$ . If  $n = 0$  choose  $v_i$  such that  $V_1 \cap V_2 = \mathbf{span} v_3$  and  $V_i = \mathbf{span} v_i + V_1 \cap V_2$  for  $i = 1, 2$ . Define the endomorphism  $a$  by  $a(v_1) = v_2$ ,  $a(v_2) = 0$ , and  $a(v_1 + v_3) = v_3$  and  $a^+$  by  $a^+(v_1) = 0$ ,  $a^+(v_2) = v_1$ , and  $a^+(v_3) = v_1 + v_3$ . Proceeding from  $n-1$  to  $n$  choose  $W$  of codimension 1 in  $V$  such that  $V_1 \cap V_2 \not\subseteq W$  and put  $W_i = W \cap V_i$ . Choose endomorphisms  $a_0, a_0^+$  of  $W$  connecting  $W_1$  and  $W_2$  according to hypothesis. Choose  $v_3 \notin W$  and  $v_i$  such  $V_i = \mathbf{span} v_i + W_i$  for  $i = 1, 2$  and extend  $a_0$  and  $a_0^+$  to obtain  $a$  and  $a^+$ , defined for  $v_i$  as above.

By this construction there are finite-dimensional  $W_n = V_n \oplus U_n$  with mutually reflexive inverses  $a_{0n}, b_{0n}$  in  $V_n$  such that  $t_{n+1}(a_{0n}, b_{0n})t_n(a_{0n}, b_{0n}) \neq t_n(a_{0n}, b_{0n})$  and isomorphism  $c_{0n} : V_n \rightarrow U_n$ . Choose  $a_n$  extending  $a_{0n}$  and  $c_{0n}^{-1}$ , and  $b_n, c_n$  extending  $b_{0n}$  and  $c_{0n}$ , respectively, such that  $b_n|_{U_n} = 0$  and  $c_n|_{U_n} = 0$ . Then the direct product of the  $\text{End}(W_n)$  provides  $R$  and  $a, b, c$  as required.

In (ii) consider a vector space  $V$  with basis  $v_n, w_n (n \in \mathbb{N})$ ,  $R = \text{End}(V)$  and define  $a(v_n) = w_n$ ,  $a(w_n) = w_{n+1}$ ,  $b(w_n) = v_n$ , and  $b(v_n) = v_{n+1}$ .  $\square$

**3.4. (Existence) varieties of unit-regular rings.** Observe that subrings of regular rings are not regular, in general, an obvious example being  $\mathbb{Z} \subset \mathbb{Q}$ . Thus, to deal with classes  $\mathcal{C}$  of regular rings in the framework of Universal Algebra, without specifying operations of quasi-inversion, it is convenient to introduce the class operator  $\mathbf{S}_\exists(\mathcal{C})$  associating with  $\mathcal{C}$  the class of all regular rings which are subrings of members of  $\mathcal{C}$ . Referring to the usual operators  $\mathbf{H}$ ,  $\mathbf{P}$ , and  $\mathbf{P}_u$  for homomorphic images, direct products and ultraproducts (which preserve regularity), a class  $\mathcal{V}$  of regular rings which is closed under  $\mathbf{H}$ ,  $\mathbf{S}_\exists$ , and  $\mathbf{P}$  (whence also  $\mathbf{P}_u$ ) is an *existence variety*, shortly  $\exists$ -variety (cf. Hall [11] for this concept). According to Herrmann and Semanova [15, Thm. 16] every existence variety of regular rings is generated by its artinian members.

For a class  $\mathcal{C}$  of regular rings let  $\mathbf{T}(\mathcal{C})$  consist of all regular rings endowed with an operation of inner inversion (that is, members of  $\mathcal{R}$  as defined in the previous subsection) where the underlying ring is in  $\mathcal{C}$ . For a ring  $R$  let  $R^{n \times n}$  denote the ring of  $n$ -by- $n$ -matrices.

- Fact 22.**
- (i) *The smallest existence variety  $\mathbf{V}_\exists(\mathcal{C})$  containing  $\mathcal{C}$  is  $\mathbf{HS}_\exists\mathbf{P}(\mathcal{C})$ .*
  - (ii)  *$R \in \mathbf{HS}_\exists\mathbf{P}_u(\mathcal{C})$  for every subdirectly irreducible  $R \in \mathbf{V}_\exists(\mathcal{C})$ .*
  - (iii) *Any subdirectly irreducible regular ring  $R$  is an  $F$ -algebra for a suitable field  $F$ . Moreover, if such  $R$  is nonartinian then  $F^{n \times n} \in \mathbf{HS}_\exists(\mathcal{R})$  for all  $n < \omega$  and  $\mathbf{V}_\exists(R) = \mathbf{V}_\exists\{F^{n \times n} \mid n < \omega\}$ .*
  - (iv) *Any identity in the language of  $\mathcal{R}$  which is valid in  $\mathbf{T}(\mathcal{C})$  is also valid in  $\mathbf{TV}_\exists(\mathcal{C})$ .*
  - (v)  *$\mathbf{TV}_\exists(\mathcal{C}) = \mathbf{VT}(\mathcal{C})$ .*

*Proof.* Referring to Herrmann and Semanova [15], (i), (iv), and (v) follow from [15, Prop. 10 (i)]. (ii) is [15, Prop. 7]. (iii) follows from [15, Thm. 16] and its proof.  $\square$

Define  $s_n(x) = t_n(x, x^+)$ . In the following, the equivalence of (5) and (9) is due, in essence, to O'Meara and Raphael [25, Thm.2.15].

**Theorem 23.** *For an existence variety  $\mathcal{V}$  of regular rings the following are equivalent (where the notion of “term” and the terms  $x^+$  and  $t_n(x)$  are as in Thm. 20)*

- (1) *All members of  $\mathcal{V}$  are perspective.*
- (2) *All members of  $\mathcal{V}$  are unit-regular.*
- (3) *All subdirectly irreducible members of  $\mathcal{V}$  are directly finite.*
- (4) *All subdirectly irreducible members of  $\mathcal{V}$  are artinian.*
- (5) *There is  $d < \omega$  such that all artinian subdirectly irreducible members of  $\mathcal{V}$  are of length  $\leq d$ .*
- (6) *There are  $d < \omega$  and a class  $\mathcal{C}$  of artinian regular rings of length  $\leq d$  such that  $\mathcal{V} = \mathbf{V}_{\exists}(\mathcal{C})$ .*
- (7) *There is  $n < \omega$  such that  $s_{n+1}(x)s_n(x) = s_n(x)$  is valid in  $\mathbf{T}(\mathcal{V})$ .*
- (8) *There is  $m < \omega$  such that the identities  $(x^{m+1})(x^{m+1})^+x^m = x^m$  and  $x^m(x^{m+1})^+x^{m+1} = x^m$  are valid in  $\mathbf{T}(\mathcal{V})$ .*
- (9) *There is a term  $u(x)$  yielding unit inner inverses, uniformly in  $\mathcal{V}$ ; that is,  $u(a)$  is a unit inner inverse for any  $R \in \mathcal{V}$  and  $a \in R$ .*

Actually, given  $d \geq 2$  in (5) one can choose  $n = d - 2$  in (7) and  $m = d$  in (8).

*Proof.* (7) implies (1) by (i) of Theorems 20. (8) implies (1), too, in view of Theorem 14. (1) is equivalent to (2) by Theorem 13, and (2) implies (3).

Each of (3) and (4) implies (5): Indeed, assume that there are artinian subdirectly irreducibles  $R_n \in \mathcal{V}$  with no bound on length. Renumbering and passing to corners and isomorphic copies, we may assume that  $R_n \cong D_n^{n \times n}$  for some division ring  $D_n$ . Thus, for fixed  $m$  and all  $n \geq m$ , the ring  $R_n$  contains a subring  $R_{mn} \cong D_n^{m \times m}$ . Choose  $R_{mn} = 0$  for  $n < m$ . Thus, in particular  $R_{mn} \in \mathcal{V}$  for all  $m, n$ . Recall that, for fixed  $m$ , the class of all rings isomorphic to  $D^{m \times m}$  for some division ring  $d$  can be finitely first-order axiomatized if one adds  $m^2$  constants for a system of matrix units. Thus, choosing a non-principal ultrafilter  $\mathcal{F}$  on  $\mathbb{N}$  one has for any fixed  $m$  the ultraproduct  $(\prod_{n \in \mathbb{N}} R_{mn})/\mathcal{F}$  isomorphic to  $D^{m \times m}$  where  $D = (\prod_{n \in \mathbb{N}} D_n)/\mathcal{F}$ . It follows  $D^{m \times m} \in \mathcal{V}$  and thus  $F^{m \times m} \in \mathcal{V}$  for all  $m$  where  $F$  is the center of  $D$ . Now, consider any infinite-dimensional  $F$ -vector space  $W$  and  $\text{End}(W_F)$ ; the latter is subdirectly irreducible, nonartinian, and not directly finite. By Fact 22(iii) one has  $\text{End}(W_F) \in \mathcal{V}$  contradicting both (3) and (4).

(5) implies (4): Assume there is subdirectly irreducible  $R \in \mathcal{V}$  which is not artinian. By Fact 22(iii),  $R$  is an  $F$ -algebra for some field  $F$  and

$\mathcal{V} \supseteq \mathbf{V}_{\exists}(R) = \mathbf{V}_{\exists}\{F^{n \times n} \mid n < \omega\}$  so that the (subdirectly irreducible)  $F^{n \times n} \in \mathcal{V}$  for all  $n < \omega$ , contradicting (5).

(5) implies (6): Since (5) implies (4), in view of Fact 22(ii) it follows that  $\mathcal{V}$  is generated by members of length  $\leq d$ .

(6) implies (7) and (8): Let  $\mathcal{R}_d$  consist of all artinian regular rings which are of length at most  $d$ . Thus,  $\mathcal{V} \subseteq \mathbf{V}_{\exists}(\mathcal{R}_d)$ . Now, consider subdirectly irreducible  $R \in \mathcal{V}$ . By Fact 22(ii) one has  $R \in \mathbf{HS}_{\exists}\mathbf{P}_u(\mathcal{R}_d)$ . Since the property of having length  $\leq d$  can be expressed, easily, by a first-order formula (in various ways), we have  $\mathbf{P}_u(\mathcal{R}_d) \subseteq \mathcal{R}_d$  while  $\mathbf{HS}_{\exists}(\mathcal{R}_d) \subseteq \mathcal{R}_d$  is obvious. This implies that  $R \in \mathcal{R}_d$  whence  $\mathcal{V} \subseteq \mathbf{V}_{\exists}(\mathcal{R}_d)$  by Fact 22(ii). By (ii) of Theorem 8 the identities  $s_{n+1}(x)s_n(x) = s_n(x)$ ,  $(x^{m+1})(x^{m+1})^+x^m = x^m$ , and  $x^m x^m (x^{m+1})^+ x^{m+1}$  (where  $n = d - 2$  and  $m = d$ ) are valid in  $\mathbf{T}(\mathcal{R}_d)$  and so in  $\mathbf{T}(\mathcal{V})$  by Fact 22(iv).

(9) implies (2), trivially. Conversely, by (v) of Fact 22, the free  $\mathbf{T}(\mathcal{V})$ -algebra  $A$  on a single generator  $a$  is unit-regular; that is, there is a term  $u(x)$  that that  $u(a)$  is a unit inner inverse of  $a$  in  $A$ . Thus  $u(b)$  is a unit inner inverse of  $R$  for all  $R \in \mathcal{V}$  and  $b \in R$   $\square$

The following has been suggested by the referee.

**Corollary 24.** *Considering classes  $\mathcal{V}$  closed under  $\mathbf{H}$ ,  $\mathbf{S}_{\exists}$ , and  $\mathbf{P}_u$  the equivalences of Theorem 23 remain valid if  $\mathbf{P}$  is replaced by  $\mathbf{P}_u$ , everywhere.*

*Proof.* For the equivalence of (1), . . . (8) it suffices to adapt (i)–(iv) of Fact 22 and to observe that  $\mathbf{P}_u\mathbf{H}(\mathcal{C}) \subseteq \mathbf{HP}_u(\mathcal{C})$  which is well known due to the fact that an ultraproduct of surjective homomorphisms amounts to a surjective homomorphism. In order to complete the proof that both (3) and (4) imply (5) observe that  $\mathbf{End}(W_F) \in \mathbf{HS}_{\exists}\mathbf{P}_u(\{F^{m \times m} \mid m < \omega\})$  by (ii) of Fact 22. and that  $F^{m \times m} \in \mathbf{S}_{\exists}\mathbf{P}_u(\{R_n \mid n < \omega\})$  whence  $\mathbf{End}(W_F) \in \mathcal{V}$ . Also (9) implies (2), trivially.

To prove that (9) implies (5) we proceed by contradiction. Assume that for each  $n$  there is  $R_n \in \mathcal{V}$  which is subdirectly irreducible artinian of length  $d_n \geq n$ . In particular, there is a division ring  $F_n$  such that  $R_n \cong M_{d_n}(F_n)$ . Let  $R$  a non-trivial ultraproduct of the  $R_n$ .  $R$  contains, for each  $n$ , a  $d_n \times d_n$  system of matrix units whence a subring isomorphic to  $M_{d_n}(F)$  where  $F$  is the prime subfield of the corresponding ultraproduct of the  $F_n$ . According to O'Meara and Raphael [25, Thm. 2.15] there is no  $u$  as required in (9).  $\square$

**Corollary 25.** *The analogues of the equivalences of Theorem 23 are valid for varieties  $\mathcal{V}$  of regular rings with inner inverse as basic operations, i.e. subvarieties of  $\mathcal{R}$  as defined in Subsection 3.3; that is, with HSP in place of  $V_{\exists}$  and omitting operator  $T$ .*

*Proof.* The only step in the proof of Thm.23 to be reconsidered is (5)  $\Rightarrow$  (4); this follows as in the proof of Cor.24.  $\square$

**Remark.** At present, only some of the implications could be extended to  $*$ -regular rings. This is related to the following questions, the first being due to Handelman. Are all  $*$ -regular rings directly finite or even unit-regular? Is the variety of all  $*$ -regular rings (with pseudo-inversion) generated by artinians? If a variety is generated by artinians does it have all members directly finite? The reasoning given in [17, 18] for a positive answer to the latter is incomplete and, most likely, cannot be completed. Thus, direct finiteness of semiartinian  $*$ -regular rings (claimed in [19]) remains open, too.

#### 4. DECLARATIONS

4.1. **Funding.** None.

4.2. **Data availability.** Not applicable.

4.3. **Ethical standards.** The author declares that there are no conflicts of interest.

#### REFERENCES

- [1] Amemiya, I., Halperin, I.: Complemented modular lattices. *Canadian J. Math.* **11**, 481–520 (1959)
- [2] Ara, P., Goodearl, K.R., Nielsen, P.P., Pardo, E., and Perera, F.: The separativity problem in terms of varieties and diagonal reduction. To appear in *Contemporary Math.*, [arxiv.org/aba/2403.11837](https://arxiv.org/abs/2403.11837)
- [3] Baccella, G., Spinosa, L.:  $K_0$  of semiartinian von Neumann regular rings. Direct finiteness versus unit-regularity. *Algebr. Represent. Theory* **20**, 1189–1213 (2017)
- [4] Bergman, G.M.: Hereditary commutative rings and centres of hereditary rings. *Proc. London Math. Soc.* **23**, 214–236 (1971)
- [5] Birkhoff, G.: *Lattice Theory*. AMS Colloq.Publ. Vol. 25, Providence RI (1967)
- [6] Ehrlich, G.: Unit-regular rings. *Portugal. Math.* **27**, 209–212 (1968)
- [7] Ehrlich, G.: Units and one-sided units in regular rings. *Trans. Amer. Math. Soc.* **216**, 81–90 (1976)
- [8] Garg, Sh., Grover, H. K., Khurana, D.: Perspective rings. *J. Algebra* **415**, 1–12 (2014)
- [9] Goodearl, K.R.: *Von Neumann Regular Rings*, second edition, Krieger, Malabar (1991)
- [10] Goodearl, K.R., Menal, P., Moncasi, L.: Free and residually artinian regular rings. *J. Algebra* **156**, 407–432 (1993)



- [11] Hall, T.E.: Identities for existence varieties of regular semigroups, Bull. Austral. Math. Soc. **40**, 59–77 (1989)
- [12] Goodearl, K.R., Menal, P.: Stable range one for rings with many units, J. Pure Appl. Alg. **54**, 261–287 (1988)
- [13] Handelman, D.: Perspectivity and cancellation in regular rings. J. Algebra **48**, 1–16 (1977)
- [14] Herrmann, C., Kindermann, M., Wille, R.: On modular lattices generated by  $1 + 2 + 2$ . Algebra Universalis **2**, 243–251 (1975)
- [15] Herrmann, C., Semenova, M.: Existence varieties of regular rings and complemented modular lattices. J. Algebra **314**, 235–251 (2007)
- [16] Herrmann, C., Semenova, M.: Linear representations of regular rings and complemented modular lattices with involution. Acta Sci. Math. (Szeged) **82** no. 3-4, 395–442 (2016)
- [17] Herrmann, C.: Direct finiteness of representable regular  $*$ -rings. Algebra Universalis **80:1** Paper No. 3, 5 pp. (2019)
- [18] Herrmann, C.: Varieties of  $*$ -regular rings. Math. Slovaca **70** no. 4, 815–820 (2020)
- [19] Herrmann, C.: Unit-regularity and representability for semiartinian  $*$ -regular rings. Arch. Math. (Brno) **56** no. 1, 43–47 (2020)
- [20] Khurana, D.: Unit-regularity of regular nilpotent elements. Algebr. Represent. Theory **19** no. 3, 641644 (2016)
- [21] Lee, T.-K.: Unit-regularity of elements in rings. J. Algebra Appl. **22** No. 6, Paper No. 2350135, 13 pp. (2023)
- [22] Mary, X.: Special clean elements, perspective elements and perspective rings. J. Algebra Appl. **22** No. 4, Paper No. 2350090, 19 pp. (2023)
- [23] Micol, F.: On Representability of  $*$ -Regular Rings and Modular Ortholattices, PhD thesis, Technische Universität Darmstadt (2003), <https://tuprints.ulb.tu-darmstadt.de/epda/000303/diss.pdf>
- [24] Nielsen, P.P., Šter, J.: Connections between unit-regularity, regularity, cleanness, and strong cleanness of elements and rings. Trans. Amer. Math. Soc. **370:3** 1759–1782 (2018)
- [25] O’Meara, K.C., Raphael, M.-R.: Uniform diagonalization of matrices over regular rings. Algebra Universalis **45** no.4, 383–405 (2001)
- [26] von Neumann, J.: Continuous Geometry. Princeton University Press, Princeton, New Jersey (1960)
- [27] Tyukavkin, D. V.: Regular rings with involution. Vestnik Moskovskogo Universiteta, Matematika **39** (1984), 29–32.
- [28] Wehrung, F.: Congruences of lattices and ideals of rings. In: Grätzer, G. Wehrung, F. (eds) Lattice theory: Special topics and applications. Vol. 1, 297–335, Birkhäuser/Springer, Cham (2014)

TECHNISCHE UNIVERSITÄT DARMSTADT FB4, SCHLOSSGARTENSTR. 7, 64289 DARMSTADT, GERMANY

*E-mail address:* herrmann@mathematik.tu-darmstadt.de