Algebra Universalis 34 (1995) 75–80

0002-5240/95/010035-29$1.50+0.20/0
© 1995 Birkhäuser Verlag, Basel

Alan Day's work on modular and Arguesian lattices

G. GRÄTZER

Dedicated to the memory of Alan Day

The present state of art in the theory of modular lattices is to a great extent due to Alan Day's contributions. The purpose of the present paper is to outline the most important ones and their impact on further developments. This is accompanied by problems of various degrees of relevance and difficulty. For the better ones, full credit should be given to Alan. For a more detailed account of some of the subjects the reader may consult Alan's excellent surveys [7, 11, 16].

1. Modular lattices with four generators

The first thing I learned from Alan about modular lattices was the following diagram drawn on the ground in the fall of 1971 — see Figure 1. The lattice is called A_{25}.

THEOREM 1. *A_{25} is the modular lattice freely generated by a, b, c, d subject to the relations*

$$ab \wedge ac = ab \wedge ad = bc \wedge bd = cd = 0$$

$$a \vee b = a \vee d = b \vee c = b \vee d = 1.$$

As it turned out, this diagram is basic for the understanding of 4-generated modular lattices. In the joint paper [12] with R. Wille the subdirectly irreducible factor R_1 of A_{25} was characterized as the free modular lattice generated by a, b, c, d satisfying the additional relation $a \vee b = 1$. Also, the interval sublattices $S[a, 1] = [a, 1]$ were

Presented by J. Sichler

Received December 7, 1992; accepted in final form October 13, 1994.



Figure 1

discovered as further subdirectly irreducibles with four generators (the doubly irreducible elements) — see Figure 2.

Freese [30] continued this investigation showing that $\mathcal{B}_2$ and its dual, the $S(5, 4)$, and the 6-element length two lattice M_4 are the only subdirectly irreducible breaches two modular lattices with four generators. In [43] and [44] the same list appeared for subdirectly irreducible modular lattices generated by two pairs of complements and for subdirectly irreducible modular 2-distributive lattices with four generators. Here, following A. Huhn [58], a modular lattice L is 2-distributive if it satisfies

$$x(y + z + t) = xy + xz + x(z + t) \cdot y(t + z).$$

Equivalent conditions are that L contains no proper 3 frame (cf. section 4) resp. no projective plane in its variety – so this covers that part of modular lattice theory where drawing of order diagrams makes sense.



Figure 2

Knowledge about 4-generated modular lattices has proved helpful with various algebraic problems. For example, a lattice theoretic version of the Baer-Kaloujnine Theorem for direct decompositions and so an alternative proof of the Krull-Remak-Schmidt theorem [40] in congruence modular varieties has been based on the list of lattices generated by two pairs of complements – and started also the investigation of permutability properties of the commutator.

Another example is the (linear) representation theory of posets resp. the free modular lattice they generate. Here a *representation* of a lattice L is a homomorphism into a vector space lattice and *indecomposable* if it does not arise as the direct product of two such. For the free modular lattice $FM(4)$ on four generators a complete list of finite dimensional indecomposable representations has been provided by Gelfand and Ponomarev [36]. The image $FM(4)$ under such a representation turned out to be D_3, M_3, M_4, one of the $S(q,4)$, or the full lattice of subspaces of a vector space over a prime field having dimension $3 \leq d \leq \infty$. And so one has a complete list of 4-generated subdirectly irreducibles in the variety generated by complemented Arguesian lattices (of [45]). The lattice J_+ occurred when considering ultraprochiers of finite dimensional representations of $FM(4)$ and this has been used for determining the *perfect elements* (i.e. those ones providing direct decompositions) of Gelfand and Ponomarev [37], Day's and Ringel [9], and [45]. The structural analysis of $FM(4)$ can even be carried over to lattices generated by posets

of *tame representation type* (for which, roughly speaking, classifying the finite dimensional representations does not involve classifying matrix pairs) see Cylke [2].

$\mathcal{A}_6$ is also behind the construction of a finitely generated modular ortholattice with an infinite set of perspective orthogonal elements by G. Bruns and M. Roddy [1].

Checking through the list of subdirectly irreducibles was the first step in the classification of pairs consisting of a finite dimensional Hermitean space E of characteristic 2 and a subspace F, see H. Gross, R. Moresi et al. [39]. Here, the lattice generators are $F, F^{\perp}, F^{*}, F^{*\perp}$ where F^{*} is the subspace of linear coherent vectors. Lattice relations on these generators allowed to single out the possible indecomposable quadruples of such subspaces and lead finally to the 12 subdirectly irreducible polarity lattices serving as the primary isometry invariants for orthogonally indecomposable pairs (E, F).

Combining examples from the list (generalized from vector spaces to abelian groups) and testing their gluing structure resulted in the proof of the unsolvability of the word problem for $FM(4)$.

Problems. Is every 4-generated Arguesian lattice embeddable into the subgroup lattice of an abelian group?

Is the word problem for free 4-generated Arguesian lattices solvable?

Is the 4-generator word problem solvable within the congruence variety of abelian groups of exponent p^2?

2. Splitting modular lattices

Following McKenzie [61] a finite subdirectly irreducible lattice L is *splitting* (within a given variety) if there is a greatest variety not containing L. For a subdirectly irreducible lattice L a sufficient condition for being splitting is to be a *bounded homomorphic image*: for every finitely generated (free) lattice M and homomorphism π of M onto L each preimage class $\pi^{-1}(x)$ has a least and greatest element. In the variety of all lattices, the splitting lattices are exactly the subdirectly irreducible bounded images [61]. An intrinsic characterization has been given by B. Jónsson and J. B. Nation [51].

Alan's results on nonmodular splitting lattices are well known. Yet, also most of what is known about splitting modular lattices can be traced back to his paper [4] cf. E. T. Schmidt [68, 69, 70].

THEOREM 2. *Let $\mathcal{V}$ be any modular lattice variety containing all 3 distributive and L a finite planar modular lattice. Then the following are equivalent:*

(1) *L is a subdirect product of splitting lattices in $\mathcal{V}$*

(2) L *is acyclic*

(3) L *is finitely projected in* $\mathcal{V}$.

Any acyclic lattice in $\mathcal{V}$ *is a bounded homomorphic image whence a subdirect product of subdirectly irreducible lattices in* $\mathcal{V}$.

Here, a finite 2-distributive modular lattice is called *acyclic* if it contains neither a sublattice M_4 nor a cycle of M_3's. The lattices in Figures 3a and 4 are acyclic, but in Figure 3b is not. Alan showed that acyclic planar lattices are *finitely projected*: for each a as above there is a finite sublattice L' of M with $\pi(L') = L$ or



Figure 3



Figure 4

equivalently, L has a finite projective cover. (The subdirectly irreducibles are their splitting, too). In this proof be considered a planar modular lattice L as a planar distributive lattice D with some new 3-irreducibles inserted in "boxes". D is finitely projective, obviously, with a planar distributive preimage D'. For a single M_3 in L one gets a preimage using the fact that Figure 4a presents the free modular lattice with generators a, b, c such that $ab < a \vee b$, $a \vee b = b$. The preimage already obtained can be adjusted since putting a chain on an edge of an M_3 yields a subdirect power as in Figure 4b. Having no cycle, this adjustment comes to an end (Figure 4c,d) — otherwise, one would spiral for ever. Finally, M_n is not a splitting modular lattice since it is a homomorphic image of R_n which is in the variety generated by $\{S_k : k \geq 3\}$ and similarly, for every planar lattice containing an interval M_n with $n \geq 4$.

Acyclic breadth 2 lattices have been studied by A. Mirchke and R. Wille [62]. In [52] acyclic lattices have been described in terms of the "geometry" on the set of join irreducibles: let a *line* be a maximal, at least 3-element set of join irreducibles such that any pair out of this set yields the same join, the *linetop*. A finite modular lattice is acyclic if and only if each of its lines is 3-element and it admits a choice of one line per linetop such that the resulting set of lines does not contain a cycle. (The irreducibles can be presented by points in the plane and the lines by segments such that the resulting figure is a union of trees). With that description, Alan's approach to the planar case could be mimicked to prove that acyclic lattices are bounded homomorphic images within the variety of modular lattices. Acyclic lattices proved important for the representation theory of modular lattices and the application to quadratic forms in infinite dimensions by M. Wild [72]. In [52] it has been shown that a finite $\mathcal{D}$-distributive modular lattice is of *finite representation type* (i.e. there are, up to isomorphisms induced by linear maps, only finitely many indecomposable representations over a fixed field) if and only if it is acyclic. And for acyclic lattices the representations (over a given field) are in 1-1 correspondence to the subdirectly irreducible factors.

The "spiral" phenomenon has been elaborated by E. T. Schmidt [67] and R. Freese [31], later on, to produce a finitely generated simple modular lattice with no prime quotients and to show that the variety of modular lattices of projective breadth 2 is not generated by its finite dimensional members.

Problems. Are acyclic breadth 2 modular lattices finitely projective?

How are the following properties of finite lattices related to each other within the variety of all modular (Arguesian) lattices: "Subdirect product of splitting lattices", "Bounded homomorphic image", "Cover bounded homomorphic image", "Finite representation type"? Are they equivalent for modular 2-dimensional lattices?

Is every finite uniquely representable lattice splitting?

Find a version of acyclicity beyond 2-dimensional!

3. Desargues' law

Beyond 2-distributivity, modular lattice theory relies heavily on some kind of (a least local) geometric point of view, for positive and negative results as well. In order to proceed to a (local) coordinatization, as in Projective Geometry one needs sufficiently high dimension or some kind of Desargues' law. Such has been provided by B. Jónsson [54, 56]: given "triangles" $\mathbf{a} = (a_0, a_1, a_2)$ and $\mathbf{b} = (b_0, b_1, b_2)$ let $c_i = (a_j \vee a_k) \wedge (b_j \vee b_k)$ for $\{i, j, k\} = \{0, 1, 2\}$ and $o = c_0 \vee (c_1 \wedge c_2)$. A lattice is *Arguesian* if it satisfies the identity

$$(o \vee b_0)((a_1 \vee b_1) \wedge (a_2 \vee b_2)) \leq a_0 \vee ((a_1 \vee o) \wedge (b_1 \vee o)).$$

This law holds in any lattice consisting of permuting equivalence relations [54] or belonging to a modular congruence variety [34]. In a more geometric version one says that a, b are *centrally perspective* at "the center of perspectivity" $o = (a_0 \vee b_0) \wedge ((a_1 \vee b_1) \wedge (a_2 \vee b_2))$ and *axially perspective* if $c_i \leq a_i + a_j$. Then, an equivalent definition is that any centrally perspective pair of triangles is also axially perspective.

The hypothesis can be made more specific: a, b form a *perspectivity configuration*, shortly PC, if $a_i \wedge o = b_i \wedge o$ for all i and if the quadruples $\mathbf{a}, o$ and $\mathbf{b}, o$ are in general position (cf. Figure 5. Here, $(x_0, \ldots, x_3)$ is in general position if $x_i \wedge (x_j \vee x_k)$ is the same $\neq 0$ for any choice of 3 distinct indices. Equivalently, $x_0 \cdots x_3$



Figure 5



Figure 9

generate a sublattice which is a subdirect product of a 3-frame generated lattice and factors D_2 where $x_i = 0$ for all but one index i—cf. Figure 6.

In a PC, one has $a_i - b_i = a_j - b_j$, $p_i p = (a_i - b_i)p(a_j - b_j)$, and (a_i, b_i, a_j, b_j) in general position for $\{i, j, k\} = \{0, 1, 2\}$. Moreover, PC's are projective configurations within the variety of modular lattices [21]. In [8] Alan showed

THEOREM 3. *A modular lattice is Arguesian if and only if every PC is axial; perspective. 2-distributive modular lattices are Arguesian.*

So one obtains the following hierarchy of modular lattice varieties, all self-dual:

distributive $\Rightarrow$ 2-distributive modular $\Rightarrow$ Arguesian $\Rightarrow$ modular.

Experience has proven that, in an axiomatic approach, Arguesian lattices or any equivalent version, cf. Alan's [25], are the proper concept: it is strong enough to allow results, and handy enough if one is going to derive them. This is more so since the important semantically defined lattice classes cannot be finitely axiomatized, see M. Haiman [42]. Behind this there is a series of higher Arguesian laws of increasing number of variables and strength (which can be separated by modified lattices of vector subspaces over any given field), each set short in variables axiomatizing consecutive varieties and modular complemented varieties (this is in [41] and R. Freese [33]).

Motivated by the same question, namely whether every Arguesian lattice has a permuting equivalence representation, Alan and St. Tsehantz [28] studied an identity due to Schützenberger and showed that it defined a lattice variety between permutable equivalence lattices and Arguesian lattices.

In another direction Alan found that in the orthomodular lattice of closed subspaces of a Hilbert space the Arguesian identity is valid for triangles consisting of orthogonal points. From this he derived the orthoarguesian identity which is satisfied by Hilbert space lattices but not in general cf. [38]. Also, he realized that orthogonal frames in Arguesian orthelattices play a central role for the investigation of varieties of modular orthelattices – which then was undertaken by M. Roddy [66].

Problems. Is there a list of finite configurations, projective within the variety of modular lattices, the exclusion of which (as relative substructures) defines the class of Arguesian lattices? Can this list be chosen consisting of enrichments of M_3's?

Is Schützenberger's identity equivalent to the Arguesian law? Is it valid in all 2-distributives?

Is there a decision procedure for the equational theory of Arguesian lattices?

Is there a finitely based modular lattice variety which is generated by its finite members and contains the subspace lattices of vector spaces over a fixed finite field?

Which representability classes of lattices form varieties? Consider e.g. vector spaces, (abelian) groups, and permuting equivalence representations – cf. [3, 43]?

Is every orthoarguesian lattice representable by means of an orthomodular space?

4. Frames and coordinate rings

A lattice theoretic concept of coordinate system has been introduced by v. Neumann [64] and in an equivalent, but sometimes more convenient form by G. Bergman and A. Hutchinson. A *n-frame* or *n-diamond* (originally called a *n-dimensional basis*) in a modular lattice L is a sequence $\mathbf{d} = (d_0, \dots, d_{n-1}) = (c_0, \dots, c_{n-1}; 0, 1)$ of elements of L such that

$$\Sigma_{j \neq i} d_j = e \qquad \text{for all } i$$

$$d_i \Sigma_{k \neq i,j} d_k = c_{ij} \qquad \text{for all } i \neq j.$$

Figure 7a and b show the correspondingmatroidal (consider only joins and those meets which satisfy the rank formula) and geometric diagrams (four points in general position and two coordinate axes with unit points and points at infinity). If



Figure 8

$a_0 = 0$, and $a_1 = 1$, one speaks of a *spanning* frame. In the standard example, L is the lattice $\mathscr{P}(\mathbb{R}^n)$ of all submodules of a free module with basis $e_0, \dots, e_{n-1}$, Re_i for $1 \le i \le n$, $e = Re_0$, and $0 = R(\Sigma_{1 \le i \le n} e_i)$. Projectively, these are the directions of the coordinate axes, the zero, and the unit point on the diagonal.

If L is the diagonal D, consisting of all complements of the point a_1 unit $e = (e - a_0)\Sigma a_i$, in the interval $[a_0, e - a_1]$ which serves as *coordinate domain* – see Figure 8. In the example, $D = \{R(a_0 + a\Sigma_{1 \le i \le n} e_i : a \in R\}$. With $b_0 = (x \mapsto x + b)$ and $b_1 = (y \mapsto y \cdot b)$, addition and multiplication are defined by

$$a \oplus b = (x \mapsto a(x - (e - a)(e - b_0)))$$

$$a \otimes b = (z \mapsto a(x \mapsto y \mapsto z(x + b \cdot y)))$$

– see Figures 9 and 10. The following general result is due to v.Neumann [64] and R. Freese [37] for the modular case, to Alan and D. Pickering [25, 8] for the Arguesian case.



Fig. 1



Fig. 2

THEOREM 4. *For an n-frame in a modular lattice L, if $n \geq 4$ or if $n \geq 3$ and L is Arguesian, then the coordinate domain D is a ring with unit under the above definitions.*

If the coordinate ring has prime characteristic p, then the sublattice generated by the frame is an $n-1$-dimensional projective geometry over the p-element field.

This and Alan's Lejy were crucial for the determination of the sublattice generated by the frame, in general, [4] and applications to the egret coil theory of

modular lattices – cf. [11]. That the coordinate ring is to a high degree independent of the particular frame chosen was shown by Alan in [12].

Problems. Determine the Arguesian lattices (or just lattices of submodules) generated by a frame and a subring of its coordinate ring – at least for some derivation rings and assuming that the subring is the full coordinate ring!

For a recursive ring, does the Arguesian lattice freely generated by a frame and a ring admit a solution of the word problem?

Analyze the relationship between the various rings associated with a skew frame and its subquotient frames in an Arguesian lattice – cf. [46]!

Is there a solution to the word problem for the modular lattice freely generated by a 3-frame?

5. Coordinatization

In their joint paper [26] Alan (see **0**, Pickering proceeded to coordinatize hyperplanes. This required some additional hypotheses, of course. They used *upper complementability* which is satisfied in the standard example: for any join g over elements from the n-frame and p with $p \leq g - 1$ there is a complement $s \leq p$ of $g - s$ (see Figure 1). A *hyperplane* is a join of n-2 elements from the frame.

THEOREM 5. *For any Arguesian lattice L with upper complementable spanning n frame, $n \geq 3$, and for every hyperplane h of such there is a lattice homomorphism of $[0, h]$ into $\mathcal{L}(_DD^n)$ having all finitely generated submodules in its image. Here D is the coordinate ring of the frame.*



Figure 1



Figure 12

Their coordinatization is the natural one in the case of planes – see Figure 12. From this they derived that L is isomorphic to the lattice $\mathcal{L}(_RD^n)$ of finitely generated submodules of $_RD^n$ with regular resp. completely primary universal D in case L is complemented resp. primary Arguesian (in a primary lattice join and meet irreducibles form trees under order and reverse order, respectively, and any interval is either a chain or has at least 3 atoms). These are, basically, the classical coordinatization results of v.Neumann [64] and Jonsson-Monk [56]. An alternative (short) proof of v.Neumann's result also came out of their analysis: consider L embedded into the subgroup lattice of an abelian group (e.g. Pöschl's embedding) and show that the canonical map from $\mathcal{L}(_RD^n)$ is an isomorphism [48].

In another direction, the coordinatization of hyperplanes was extended by McGetrick [38] to Arguesian lattices with an additional geometric structure tailored for submodule lattices, see [71].

Problems. Is there a common generalization of the coordinatization of complemented and primary lattices which also has the potential of computing subalgebras?

For a modular lattice with spanning n-frame, $n \geq 4$, and hyperplane b, is the interval $[0, b]$ Arguesian or even embeddable into $L(D^{n-1})$?

Has every Arguesian lattice with spanning n-frame, $n \geq 3$, a permutative equivalence representation?

6. Quasiplanes

From an equational point of view, projective planes may be seen as modular lattices of primitive length ≤ 3 or as 3-distributive lattices. In the first approach, the



FIGURE 13

model class is close to the intended one but, working with the equations becomes rather awkward. The second approach includes too many models much more complicated than projective planes. Therefore, Alan [6, 7, 13] suggested to understand them as *quasiplanes* – modular lattices not containing any 2-dimensional gluing of nondegenerate 3-frames of Figure 13.

Here, two n-frames $\mathfrak{a}$ and $\mathfrak{b}$ in a modular lattice form a n-1-dimensional gluing, if $\Sigma_{\mathfrak{a}} a_{n-1}$ transposes up to $\Sigma_{\mathfrak{b}} b_1$ the matching the canonically induced n-1-frames (see [13, 1], for the generalization to k-dimensional gluings of an n and an m-frame). Gluings of non-degenerate frames can be understood as configurations projective within modular lattices, so their exclusion defines a variety. This variety can be also defined via a handy distributivity condition relatively to nondegenerate frames.

$$L \text{ modular 2-distributive} \;\Rightarrow\; L \text{ quasiplane} \;\Rightarrow\; L \text{ modular 3-distributive.}$$

Relatively complemented modular 3-distributive lattices are quasiplanes. Alan's central result [8, 6] shows how to characterize projective planes.

THEOREM 6. *For a modular lattice L the following are equivalent:*
(1) *L is an irreducible projective plane.*

(2) L *is a subdirectly irreducible quasiplane generated by a 3-frame and its coordinate domain*

(3) L *has a spanning 3-frame with respect to which all nonzeros are invertible.*

Here, the coordinate domain of a 3-frame in a modular lattice is endowed with a ternary operation from which an additive loop and a multiplication with unit can be derived [26]. In the case of projective planes this yields a ternary ring, indeed.

Problems. Is a subdirectly irreducible quasiplane generated by a 3-frame and a subset of its coordinate domain necessarily a projective plane?

Is the word problem for (free) quasiplanes (or modular 2-distributives) in more than 4 generators decidable?

Characterize quasiplanes in the geometric language for spatial modular lattices [51]!

Do all Arguesian quasiplanes admit a permutable equivalence representation?

Is every quasiplane (modular 2-distributive) variety of finite height generated by a finite lattice?

Let L a finitely generated modular lattice of finite length such that every of its complemented intervals is locally finite. Is L finite (at least if it is a quasiplane)? The answer is "yes" for 2-distributives.

7. Properties of rings

The example of embedding the lattice of $n-3$ dimensional space over the quaternions in that of a 6-dimensional space over the complex numbers shows that commutativity of the coordinatizing ring cannot be equationally expressed without a dimension restriction. In [6] Alan studied some possible approaches, each based on a projective configuration. A *strong link pair* consists of a_0, a_1, b_0, b_1 subject to relations which, in a modular lattice, grant that the sublattice generated is a subdirect product of a factor where these form a 3-frame and a factor where $a_0 = a_1$ and $b_0 = b_1$, cf. Figure 14.

A modular lattice is (SLP-) *Pappian* if for each strong link pair and $a_2 < a_0 \cdot a_1, b_2 < b_0 \cdot b_1$ one has

$$(a_0 \cdot b_1 + a_2 \cdot b_0)(a_0 \cdot a_1 + b_2)(b_0 \cdot b_1 + a_2)$$
$$< (a_0 + b_2)(a_2 + b_0) + (a_1 + b_2)(a_2 + b_1).$$

See Figure 15 where x denotes the left hand side. This, of course, reflects Pappus' Law from Projective Geometry. The dimension restriction implied by this identity



Figure 14



Figure 15

is 3-distributivity. Requiring a frame, even one with 3-distributivity (which cannot be improved in view of the above example).

THEOREM 7. *R is strongly regular if and only if $\mathcal{L}(_RR)$ is a quasiplace.*

If $\mathcal{L}(_RR)$ is Peppian then R is commutative. For regular R the converse is also true.

Here, a ring is strongly regular if for every a there is x with $a^2x=a$ or, equivalently, it is regular and $a^2=0$ only for $a=0$. These rings are of interest since they allow a distant sheaf representation.

Problem. Are there, besides the above and divisibility conditions for integers, any other properties of rings which can be expressed via identities for the lattice $\mathcal{N}_l(R)$? How about commutativity and regularity?

Is every Pappian lattice Arguesian?

Is every finite Arguesian quasiprime Pappian?

8. Critical configurations for the Arguesian identity

In a series of joint papers with B. Jónsson [20, 21, 22, 23] Alan took up a theme from his earlier work [9]. How to understand the failure of the Arguesian law in terms of well described forbidden configurations, if not orbeticules? The substantial approach turned out invisible since D. Pickering [65] found later on an infinite list of finite minimal non Arguesian modular lattices and even a modular non-Arguesian variety whose finite members all are Arguesian.

So, what they did is to study non-Arguesian (i.e. $c_2 \not\leq c_0 + c_1$) PC's and to gain as much information as possible about them. Here, the p_i, c_i, $a_i b_i$ are also included into the concept of a PC. They undertook an intricate analysis of how to construct new non-Arguesian PC's from given ones aiming at *primes* where $c_2 \leq c_0 + c_1$ or $a_i a_j$. More conceptually, such a PC has a specific *atom cover* in the pointwise ordered set of PC's.

THEOREM 3. *Every modular non-Arguesian lattice contains a prime PC or lies in its ideal lattice.*

For a prime PC in a modular lattice L there are order preserving maps α, β (depend in terms of the PC) of 2^3 into L, such that $I_\varepsilon = [\alpha\varepsilon, \beta\varepsilon]$ is a projective geometry of dimension at most 3 for $\varepsilon = 0, 1$, an irreducible projective plane otherwise. The union of these intervals contains the PC and the $p_i, c_i, a_i b_i$.

If the PC is stable and Boolean then α and β factor through ε over $2^3_{\varepsilon \neq 3}$, and $\bigcup_\varepsilon I_\varepsilon$ is a subalgebra.

Here, the PC is *Boolean* if α and β are lattice homomorphisms. Of course, some of the intervals I_ε may coincide. If that happens for all transposed $I_\varepsilon, I_{\varepsilon'}$ with $\varepsilon \wedge \varepsilon' = 0$ then the PC is called *stable*.

From each prime non stable PC one obtains a stable and Boolean one by passing down in the order of PC's. Examples for any $r \geq 3$ can be found in the next section.

On the other hand, for a stable non Boolean prime PC all of the twenty planes I_ε, ε of rank 2 or 3, are distinct. An example where this occurs was first constructed in [19], see Figure 16.



Figure 16

Problems. Are there stable prime PC's with 22 or more intervals more complicated than any other?

Can the Arguesian law be decently expressed in the geometric language for spatial modular lattices [5]?

Is there a configuration of join irreducibles (points) and meet irreducibles (hyperplanes), of finitely many such, the exclusion of which characterizes Arguesian lattices among finite length modular lattices?

9. Arguesian lattices and gluing

The above suggests to consider how the gluing construction and the Arguesian law interact. Alan, of course, had his own categorical view of gluing, as did Jónsson [16] resulting in a more general and more accessible presentation. Analytically, the most convenient way is to think in terms of "skeleton" lattices S and join resp. meet

preserving 1-1 maps σ, π of S into L with $\sigma x < \pi x$ such that L is covered by the blocks $I_x = [\sigma x, \pi x]$. If necessary, one resorts to "imaginary" bounds in a modular ideal containing both all ideals and filters of L. The tolerance γ on S indicates whether for $x \leq_S y$ the blocks overlap, $x \leq_\gamma y$ iff $\sigma y \leq \pi x$. One speaks of a *gluing* if the equivalence relation generated by γ is total.

Synthetically, given S, γ, the blocks I_x and a pair of adjunctions between each I_x, I_y, $x <_\gamma y$ one can form the disjoint union with the order defined by that on the blocks and the adjunctions (a variant of Płonka sum). Then, provided appropriate axioms are satisfied, one obtains the glued sum (with overlaps as encoded in γ) as homomorphic image. The gluing of modular lattices is always modular.

For Arguesian lattices, the situation is much more complicated. Still, Alan's papers [18, 23] provide some sufficient conditions (for the generalization one just has to notice that each complemented interval is contained in some block). Call a non-Arguesian PC *minimal* if it is so in the pointwise order of PC's.

THEOREM 9. *Let L be a gluing of Arguesian lattices. If S is a chain, or S is of length or width 2, or S is of width ≤ 3 then every minimal non-Arguesian PC is Boolean with 16 or more 2, 4 and 8 intervals, respectively.*

On the other hand, L is Arguesian provided that $S = 2$ and $I_0 \to I_1$ is distributive or, else, S is modular of finite length and projective breadth ≤ 2 and $\{I_x : x \in S\}$ is irredundant for each subdirect $T \subseteq 2^S$, $0 < 0_T$ of S.

Problems. Is there a non-Boolean minimal non-Arguesian PC in a finite length modular lattice L with $S(L)$ modular and/or 3-distributive?

In the second half of the Theorem, are the cases $n = 4, 5$ to be considered; is it sufficient to have S modular and/or 3-distributive?

Is there a non-3-distributive Arguesian lattice variety closed under gluings with $S = 2$? According to D. Pickering [65], lattices in such a variety have characteristic 2, since whereas the variety cannot be closed under gluings with S the projective plane of order 2.

10. Arguesian lattices of finite length and representability

If L is modular of finite length, then one has a particular gluing, the blocks of which are the maximal complemented intervals. S is then also addressed to as the *prime skeleton* $S(L)$.

Using the gluing of Arguesian blocks, Alan and his coauthors produced in [19, 23] typical and partly also minimal, non-Arguesian lattices illustrating the theory of the preceding sections. These examples also give some insight into conditions granting representability.



Figure 17

The basic observation dates back to B. Jónsson [55]; in essence: Frames glued 2-dimensionally in an Arguesian lattice have isomorphic coordinate rings. If the blocks are coordinatized as in the standard example then L is Arguesian (and embeddable into a submodule lattice over the coordinate ring) if and only if the gluing maps are induced by a semilinear map.

For L of length ≤ 5, by the preceding theorem, every minimal non-Arguesian PC has at most 4 (proper) covers. Accordingly, assume that $S(L)$ is an M_n, $n \geq 2$, and that for any pair of neighbours we have Arguesian 2-dimensional gluings of irreducible projective planes – cf. Figure 17.

Then the gluing maps are induced by division ring isomorphisms. Under these hypotheses, L is Arguesian (and representable over a division ring) if and only if the composition of the isomorphisms along a gluing path depends on start and goal only – cf. G. Kurlinai [60]. As a consequence of this and B. Jónsson [54, 55], Alan and his coauthors derived (cf. [33]):

THEOREM 10. *A finite length modular lattice L of length ≤ 5 or with $S(L)$ a chain is Arguesian if and only if it has a geometric coordinatization representation.*

In contrast, an Arguesian lattice of length 9 or with $S(L)$ distributive is not necessarily representable – this is witnessed by examples due to M. Hall-an [42].

For L of length 6, if L is not Arguesian, then either there is a prime Boolean PC or $S(L)$ is an irreducible plane and blocks are pairwise isomorphic planes. The first kind of example, with $S(L) = 2^n$, appeared in D. Pickering [65] (cf. Fig. 18) and was generalized there to produce his infinite list of minimal non-Arguesian lattices. The second can be also chosen as a minimal non-Arguesian lattice which, moreover, is primary and so a projective (Jónsson class; cf. [27] for more on such.



Figure 18

Problems. Has every Arguesian lattice of length 6 (7) a permuting equivalence representation?

Can representability be understood in terms of gluing of coordinatizable parts? On this there are some unpublished results of M. Haiman and J. B. Nation.

For finite lattices, is representability a decidable property?

For fixed n, is the class of representable lattices of length $\leq n$ finitely axiomatizable? Again, consider your favorite representability.

11. Databases

Alan was among the first to realize the connection of lattice theory to dependencies in relational databases. In [14] he gave a detailed analysis of the relationship between functional dependencies and sectional lattice congruences. As he pointed out, modular lattice theory naturally comes in when studying embedded multivalued dependencies. From a mathematical point of view, a database is a family $(\theta_A)_{A\in\mathcal{A}}$ of partitions on a set, each partition named with an "attribute". The underlying set can be seen as the set of rows and two rows having the same value of a given attribute will be in the same class of the partition. A functional dependency $X \to Y$ now amounts to $\theta_Y \equiv \theta_X$ where $\theta_X = \bigcap_{A\in X}\theta_A$. Following Alan [15], an embedded multivalued or symmetric dependency $[X, Y]$ is satisfied if $\theta_X \circ \theta_Y = \theta_X \vee \theta_Y$. A multivalued dependency is one where $X \cup Y = U$, the total set of attributes. An *implication* $\Sigma \to \sigma$ between a finite set of dependencies and a dependency is (finitely) valid if σ is satisfied in all (finite) databases satisfying Σ. An important

question for each kind of dependencies is to find a sound and complete axiomatization of this semantic consequence operator or an algorithm deciding whether an implication is (finitely) valid: the (finite) *implication problem*, see the survey of P. Kanellakis [59]. Positive and negative results have been obtained in various cases.

On the basis of what happens in terms of meets and products of equivalence relations, Alan suggested in [15] a *natural lattice interpretation* of dependencies via sets of relations: A map $\varphi : U \to L$ is a model for $X \to Y$ if $\Pi_\varphi X \leq \Pi_\varphi Y$ and a model for $[X, Y]$ if for all $Z \subseteq U \setminus (X \cup Y)$ one has $\Pi_\varphi(Z \cup (X \cap Y)) = \Pi_\varphi(Z \cup X) \cap \Pi_\varphi(Z \cup Y)$. So, with an implication of dependencies there corresponds a conjunction of universal Horn lattice formulas. A lattice variety $\mathcal{V}$ provides a *sound and complete* natural interpretation if this conjunction is valid in $\mathcal{V}$ if and only if the implication is valid for databases. Alan showed in [13]:

THEOREM 11. *For the class of functional and multivalued dependencies, every non-trivial lattice variety provides a sound and complete natural interpretation — and thus yields a new solution of the (finite) implication problem.*

For embedded multivalued dependencies such interpretation fails to be in terms of a non-distributive modular variety.

In particular, the rather effective methods of solving word problems in the class of all lattices can be applied. But the following remains open:

Problem. Is there a non-distributive modular lattice variety providing a sound and complete (natural) interpretation of embedded multivalued dependencies? Does such have to be Arguesian?

Though, in the same mood one sees that the (finite) implication problem is unsolvable for embedded multivalued dependencies [50]: given an n-frame, $n \geq 3$, in a partition lattice and finitely many elements of the coordinate domain plus some auxiliary ones, if one requires sufficiently, but finitely many permutability and meet relations between these partitions, then they generate a sublattice of the partition lattice which is a sublattice of the congruence lattice of a suitably chosen module [48]. Since meet and permutability relations can be captured by embedded multivalued dependencies (it needs a little trick of Vardi's to eliminate functional dependencies), one can interpret an unsolvable word problem for semigroups into implications.

12. Frames in congruence modular varieties

The ring showing up in the above had been identified by Alan in a fine paper with E. Kiss [24], earlier. There they clarified the relationships between various

rings arising in Co-ordinatization and Commutator Theory – establishing one o the essential links between Lattice Theory and Universal Algebra.

THEOREM 12. *For a n-frame, $n \geq 3$, with top τ and bottom β, of congruences of an algebra A in a congruence modular variety $\mathcal{V}$, the following rings are canonically isomorphic:*

(1) *The n-Stone – n-Newman projective coordinate ring.*

(2) *The Day–Pickering coordinate ring.*

(3) *The subdirect product of the Euclid–Hilbert affine coordinate rings of the classes of τ/β.*

(4) *The Jónsson ring of equivalence relations on A/β derived from any of the associated 2-frames.*

(5) *The Day–Kiss ring of equivalence classes of endomorphisms of the pair $(A/\beta, \tau/\beta)$.*

(6) *The Pierce–McKenzie ring $R(\mathcal{V})$ of the variety, provided the frame is free in the variety.*

If the variety $\mathcal{V}$ is residually small and either solvable or locally finite then a congruence variety is that of $R(\mathcal{V})$-modules in distributive.

Problems. Has the congruence variety of groups a decidable equational theory? Is it generated by its finite members, even?

REFERENCES

[1] Birkhoff, G. and Kiss, E., *A ternary generation of modular semilattices*, Period. Math. Hung. 27 (1993), 25–31.

[2] Chajda, I., *Polars and linearly equational classes of modular lattices*, Proceedings of the Conf. Representations of Algebras, Ottawa, 1992, to appear in Proc. Int. Conf. Representations of Algebras, Ottawa, 1992, Can. Math. Soc. Conf. Proc.

[3] Czédli, G. and Horváth, E., *Congruence distributive varieties*, Acta Sci. Math. (Szeged) 57 (1993), 35–45.

[4] Day, A., *Splitting algebras and a weak notion of projectivity*, Algebra Universalis 5 (1975), 153–162.

[5] Day, A., *Equational theories of projective geometries*, Contributions to Lattice Theory, Coll. Math. Soc. J. Bolyai, Szeged, vol. 33, 1980, pp. 277–318.

[6] Day, A., *A remark on a Pappian Moufang identity*, Canadian Math. Bull. 24 (1981), 107–108.

[7] Day, A., *Geometrical applications in modular lattices*, Universal Algebra and Lattice Theory (Proc. Conf. Charleston, S.C., 1982), Lecture Notes in Mathematics, vol. 1004, Springer-Verlag, New York, 1983, pp. 127–141.

[8] Day, A., *A note on projective geometries in modular and Arguesian lattices*, Canadian Math. Bull. 26 (1983), 283–290.

[9] Day, A., *A note on Arguesian lattices*, Arch. Math. (Basel) 49 (1987), 117–123.

[10] Day, A., *On some geometric properties defining classes of rings and varieties of modular lattices*, Algebra Universalis 27 (1990), 21–34.

[11] Day, A., *Applications of commutator theory in modular lattices: theorems of J. von Neumann*, Order 7 (1990), 395–398.

[2] DAY, A., *Some characterization results in the characterization of Arguesian lattices*, Universal Algebra and Lattice Theory, S. Comer, ed., Lecture Notes in Math. 1149, Springer-Verlag, New York 1985, pp. 37–45.

[3] DAY, A., *Characterizations of modular lattices*, Algebra Univ. 23 (1986), 1–6.

[4] DAY, A., *The lattice theory of functional dependencies and normal decompositions*, Int. J. Algebra and Computation 2 (1992), 409–431.

[5] DAY, A., *A note on the congruence extension property*, Algebra Universalis 1 (1971), 234–235.

[6] DAY, A. and HERRMANN, C., *Gluings of modular lattices*, Order 5 (1988), 85–101.

[7] DAY, A., HERRMANN, C., JÓNSSON, B., NATTRASS, D. and PICKERING, D., *Small non-arguesian lattices*, Algebra Universalis 31 (1994), 69–94.

[8] DAY, A. and JÓNSSON, B., *The structure of non-Arguesian lattices*, Bull. Amer. Math. Soc. 74 (1968), 127–129.

[9] DAY, A. and JÓNSSON, B., *A structure characterization of finite Arguesian lattices*, Order 5 (1988), 333–350.

[10] DAY, A. and JÓNSSON, B., *Non-Arguesian configurations in modular lattices*, Acta Sci. Math. (Szeged) 51 (1987), 309–318.

[11] DAY, A. and JÓNSSON, B., *Non-Arguesian configurations and pairings of modular lattices*, Algebra Universalis 26 (1989), 208–215.

[12] DAY, A. and KISS, E., *Frames and duals of continuous modular varieties*, Journal of Algebra 109 (1987), 479–507.

[13] DAY, A. and PICKERING, D., *A note on the Arguesian lattice identity*, Proc. Vanderbilt Conference on Universal Algebra (1982).

[14] DAY, A. and WILLE, R., *The characterization of Arguesian varieties*, Trans. Amer. Math. Soc. 278 (1983), 307–322.

[15] DAY, A. and STEPHAN, J., *Congruence relations on lowness semilattices*, preprint (1990).

[16] DAY, A. and TSCHANTZ, S., *Lowlies algebras, universal enveloping algebras and the Arguesian law*, preprint (1990).

[17] DEAN, V. and RINGEL, C. M., *Perfect elements in free modular lattices*, Math. Ann. 267 (1984), 97–101.

[18] DILWORTH, R., *Dependence relations in a semi-modular lattice*, Duke Math. J. 11 (1944), 575–587.

[19] DILWORTH, R., *Some structures of modular lattices not generated by their finite dimensional members*, in Coll. Math. Soc. J. Bolyai 17, Contributions to General Algebra, B. Csákány and E. Schmidt, eds., North-Holland Publ. Co., Amsterdam, 1977, pp. 111–116.

[20] DILWORTH, R., *Projective geometries as projective modular lattices*, Trans. Amer. Math. Soc. 251 (1979), 329–343.

[21] DILWORTH, R., *Finitely based modular varieties are finitely generated*, Algebra Universalis 1 (1971), 204–214.

[22] DREESE, R. and JÓNSSON, B., *Congruence modularity implies the Arguesian identity*, Algebra Universalis 6 (1976), 225–228.

[23] FREESE, R. and MCKENZIE, R., *Commutator Theory for Congruence Modular Varieties*, London Math. Soc. LN 125, Cambridge, 1987.

[24] GELFAND, I. M. and PONOMAREV, V. A., *Problems of linear algebra and classification of quadruples of subspaces in a finite dimensional vector space*, Coll. Math. Soc. J. Bolyai 5, *Hilbert Space Operators*, Tihany 1970, North-Holland, Amsterdam, 1972, 163–237.

[25] GELFAND, I. M. and PONOMAREV, V. A., *Arrangements of linear subspaces and their representations*, Russian Math. Surveys 29 (1974), 1–56.

[26] GRÄTZER, G., *Two Mal'cev-type theorems in universal algebra*, J. Combinatorial Theory 8 (1970), 334–342.

[39] GROTE, H., HEITMANN, C. and MEYER, R., *The interpretation of subspaces of Arguesian lattices*, J. Alg. 89 (1984), 516–541.
[40] GUMM, H. P. and HERRMANN, C., *Algebras in modular varieties: Tame congruences and commutators*, Houston J. Math. 7 (1981), 503–521.
[41] HAGEMANN, V., *Proof theory of finite lattices*, Arch. f. Math. 45 (1985), 289–343.
[42] HAGEMANN, V., *Equational bases which are not finitely based*, Algebra Universalis 18 (1984), 178–181.
[43] HERRMANN, C., *On modular lattices generated by two complemented pairs*, Houston J. Math. 10 (1984), 511–529.
[44] HERRMANN, C., *A criterion for subdirectly irreducible modular lattices with four generators*, Acta Sci. Math. 47 (1984), 169–175.
[45] HERRMANN, C., *Relations and subgroups*, Colloquium on modularity, Algebra Universalis 14 (1982), 207–209.
[46] HERRMANN, C., *On the word problem for modular lattices with four free generators*, Math. Ann. 201 (1973), 313–332.
[47] HERRMANN, C., *On the equivalence of projective covering systems*, Trans. Amer. Math. Soc. 235 (1978), 75–85.
[48] HERRMANN, C., *Finite sets of projective geometries*, Acta Sci. Math. 37 (1975), 99–107.
[49] HERRMANN, C., *On the representation of modular lattices*, Colloq. Math. 49 (1984), 275–281.
[50] HERRMANN, C., *On the undecidability of implications in three-generated modularity*, Universal algebra and its links with logic, algebra, combinatorics and computer science, Darmstadt 1983, Heldermann Verlag, Berlin, 1984, pp. 139–159.
[51] HERRMANN, C., PICKERT, G. and ROGGEV, M., *Geometric description of modular lattices*, Algebra Universalis 17 (1984), 301–326.
[52] HERRMANN, C. and WILD, M., *Algebraic modular lattices and their representations*, J. Algebra 85 (1983), 57–86.
[53] HIGGS, D. A., *Two-valued sentential logics*, Coll. Math. Soc. J. Bolyai 33, Universal Algebra, B. Csákány, E. T. Schmidt and J. Szendrei (eds.), North-Holland, Amsterdam, 1982, pp. 337–343.
[54] JÓNSSON, B., *On the representation of lattices*, Math. Scand. 1 (1953), 193–206.
[55] JÓNSSON, B., *Arguesian lattices of dimension n ≤ 4*, Math. Scand. 7 (1959), 133–145.
[56] JÓNSSON, B. and MONK, G., *Representations of primary Arguesian lattices*, Pacific J. Math. 30 (1969), 95–139.
[57] JÓNSSON, B. and NELSON, J. E., *A report on minimal splittings of a free lattice*, Coll. Math. Soc. J. Bolyai 17, Contributions to Universal Algebra, B. Csákány and J. Szendrei (eds.), North-Holland, Amsterdam, 1977, pp. 225–257.
[58] KAMMACHER, G., *Orthocomplete Lattices*, Academic Press, London, 1965.
[59] KANELLAKIS, P. C., *Elements of relational database theory*, in *Handbook of Theoretical Computer Science*, vol. B, *Formal Methods and Semantics*, J. van Leeuwen ed., Elsevier, Amsterdam, 1990, pp. 1073–1156.
[60] KURINOV, G. I., *A necessary condition for the lattice embeddability of finite dimensional modular lattices into complemented modular lattices* (Russian), Ordered sets and lattices (Saratov University) 3 (1978), 64–70.
[61] MCKENZIE, R., *Equational bases and non-modular lattice varieties*, Trans. Amer. Math. Soc. 174 (1972), 1–43.
[62] MEDVEDEV, A. and WILLE, R., *Über freie modulare Verbände FM(3,3)*, in *Contributions to General Algebra*, H. Kaiserslautern et al. eds., Vienna, J. Heyn, Klagenfurt, 1979, pp. 219–233.
[63] NATAN, J. B. and PICKERING, D., *Arguesian lattices whose skeleton is a chain*, Algebra Universalis 24 (1987), 81–110.
[64] v. NEUMANN, J., *Continuous Geometry*, Princeton Univ. Press, Princeton, 1960.
[65] PICKERING, D., *Minimal non-Arguesian lattices*, Ph.D. thesis, Univ. of Hawaii, 1980.
[66] ROGGEV, M., *Varieties of modular orthoalgebras*, Order 4 (1987), 405–415.
[67] SCHMIDT, E. T., *On finitely generated regular modular lattices*, Per. Math. Hung. 16 (1985), 313–316.
[68] SCHMIDT, E. T., *On splitting modular lattices*, Coll. Math. Soc. J. Bolyai 33, Universal Algebra, North-Holland, Amsterdam, pp. 697–703.

[6] SCHIEF, W. K., *Classical and integrable projective surfaces: variational principles and Gauss–Codazzi systems*, Int. Math. Res. Not. 2003, no. 71, 3809–3843.

[7] SCHIEF, W. K., *On the unification of classical and novel integrable surfaces. II. Difference geometry*, Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. 459 (2003), no. 2039, 1857–1877.

[8] SCHIEF, W. K., *Isothermic surfaces in spaces of arbitrary dimension: integrability, discrete geometry, and symmetry transformations*, Acta Appl. Math. 88 (2005), no. 2, 129–146.

[9] SCHIEF, W. K., *Moutard and Calapso–Guichard nets: projective geometry and integrability*, J. Geom. Phys. 56 (2006), no. 9, 1618–1649.

[10] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[11] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[12] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[13] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[14] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[15] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[16] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[17] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[18] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[19] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

[20] SCHIEF, W. K., *Isothermic surfaces in spaces of higher dimension: Christoffel duality, Darboux points and integrability*, Oberwolfach Preprint, OWP 2006-08, 2006.

Technische Universität Darmstadt
Fachbereich Mathematik
Schloßgartenstraße 7
D-64289 Darmstadt
Germany