

RAMSEY'S THEOREM FOR PAIRS AND PROVABLY RECURSIVE FUNCTIONS

ALEXANDER KREUZER AND ULRICH KOHLENBACH

ABSTRACT. This paper addresses the strength of Ramsey's theorem for pairs (RT_2^2) over a weak base theory from the perspective of 'proof mining'. Let RT_2^{2-} denote Ramsey's theorem for pairs where the coloring is given by an explicit term involving only numeric variables. We add this principle to a weak base theory that includes weak König's lemma and a substantial amount of Σ_1^0 -induction (enough to prove the totality of all primitive recursive functions but not of all primitive recursive functionals). In the resulting theory we show the extractability of primitive recursive programs and uniform bounds from proofs of $\forall\exists$ -theorems.

There are two components this work. The first component is a general proof-theoretic result, due to the second author ([13, 14]), that establishes conservation results for restricted principles of choice and comprehension over primitive recursive arithmetic PRA as well as a method for the extraction of primitive recursive bounds from proofs based on such principles. The second component is the main novelty of the paper: it is shown that a proof of Ramsey's theorem due to Erdős and Rado can be formalized using these restricted principles.

So from the perspective of proof unwinding the computational content of concrete proofs based on RT_2^2 the computational complexity will, in most practical cases, not go beyond primitive recursive complexity. This even is the case when the theorem to be proved has function parameters f and the proof uses instances of RT_2^2 that are primitive recursive in f .

1. INTRODUCTION

Ramsey's theorem for pairs and two colors RT_2^2 has been at the center of a lot of research in computability theory and reverse mathematics aiming at determining the complexity of the homogeneous sets in RT_2^2 and the contribution to the provably recursive functions of RT_2^2 when added to theories such as RCA_0 from reverse mathematics (see e.g. [20, 9, 8, 18, 3, 7, 19, 6]). One of the main open questions (see [3]) is whether the provably recursive functions of $RCA_0 + RT_2^2$ are the primitive recursive ones or whether the totality of the Ackermann function can be established in this system. From the perspective of applied proof theory ('proof mining') this

2000 *Mathematics Subject Classification.* 03F10; 03F35; 05D10.

This main results of this paper are from the Diploma Thesis [16] of the first author written under the supervision of the second author.

The authors gratefully acknowledge the support by the German Science Foundation (DFG Project KO 1737/5-1).

question is of relevance for determining what type of bounds one can expect to be extractable from concrete mathematical proofs of - say - Π_2^0 -sentences $\forall m \in \mathbb{N} \exists n \in \mathbb{N} A_{qf}(m, n)$ or sentences $\forall f \in \mathbb{N}^{\mathbb{N}} \exists n \in \mathbb{N} A_{qf}(f, n)$ (with A_{qf} quantifier-free) that are based on RT_2^2 . Experience from the logical analysis of many proofs in different areas of mathematics indicates that, typically, proofs of theorems $\forall f \in \mathbb{N}^{\mathbb{N}} \exists n \in \mathbb{N} A_{qf}(f, n)$ that make use of second order principles $\forall g P(g)$ such as RT_2^2 that state that for all functions g or sets of a certain type some property (here for all colorings c a property $\text{RT}_2^2(c)$) holds only need explicit instances $\psi(f)$ for g resp. c that are effectively definable in the parameter f by some closed term ψ of the underlying system \mathcal{T} , i.e.

$$\mathcal{T} \vdash \forall f \in \mathbb{N}^{\mathbb{N}} (\text{RT}_2^2(\psi(f)) \rightarrow \exists n \in \mathbb{N} A_{qf}(f, n)).$$

In this paper we show, that in such a situation and for sufficiently weak systems \mathcal{T} the extractability of a primitive recursive functional Φ (in the sense of Kleene) with

$$\forall f A_{qf}(f, \Phi(f))$$

is guaranteed. Moreover, the proof theoretic method used provides an extraction algorithm for Φ from a given proof.

We work in a setting based on fragments of (extensional) arithmetic formulated in the language of functionals of all finite types. In [10] (see also [15]), the second author introduced a hierarchy $\text{E-G}_n\text{A}^\omega$ of such fragments containing functionals corresponding to the n -th level of the Grzegorzczuk hierarchy and quantifier-free induction.

As usual in proof mining, universal axioms do not matter and so arbitrary true (in the sense of the full set-theoretic type structure over \mathbb{N} , see [15]) universal sentences can always be added to the theories used in our paper.¹

The union of all these systems is denoted by $\text{E-G}_\infty\text{A}^\omega$ and contains terms for all primitive recursive functions but not for all primitive recursive functionals (in the sense of Kleene) of type level 2 (e.g. not $\Phi_{it}(f, x, y) := f^{(x)}(y)$). This distinguishes the system from $\widehat{\text{E-PA}}^\omega \upharpoonright$ in [10, 15] (sometimes also denoted by PRA^ω).

As the theory \mathcal{T} in the result above we may take

$$\mathcal{T} := \text{E-G}_\infty\text{A}^\omega + \text{QF-AC} + \text{WKL},$$

where QF-AC is the union of the schemata of quantifier-free choice from functions to numbers

$$\forall f \in \mathbb{N}^{\mathbb{N}} \exists n \in \mathbb{N} A_{qf}(f, n) \rightarrow \exists F \forall f \in \mathbb{N}^{\mathbb{N}} A_{qf}(f, F(f))$$

¹In [10] we officially added all true universal sentences as axioms. As in the convention made in chapter 13 of [15], we in this paper instead only add universal sentences that are provable in $\widehat{\text{E-PA}}^\omega \upharpoonright$, see below) which covers, in particular, the schema of quantifier-free induction. In this way we can state various conservation results over primitive recursive arithmetic PRA but still can add further universal axioms as might be useful in concrete proofs.

and quantifier-free choice from numbers to functions

$$\forall n \in \mathbb{N} \exists f \in \mathbb{N}^{\mathbb{N}} A_{qf}(n, f) \rightarrow \exists F \forall n \in \mathbb{N} A_{qf}(n, F(n)) \quad (A_{qf} \text{ quantifier-free})$$

and WKL is the weak König's lemma (i.e. König's lemma for 0/1-trees, see [19, 15]).

It is clear that \mathcal{T} contains WKL_0^* (defined in [19]) via the usual embedding.

For this system, the second author has shown in [13, 14] that the addition of the use of fixed instances $\Pi_1^0\text{-CA}(\varphi(f))$ of Π_1^0 -comprehension

$$\Pi_1^0\text{-CA}(f) := \exists g \in \mathbb{N}^{\mathbb{N}} \forall x \in \mathbb{N} (g(x) = 0 \leftrightarrow \forall y \in \mathbb{N} (f(x, y) = 0))$$

only causes primitive recursive provably recursive functions. More precisely, by the proof of corollaries 4.4 and 4.5 in [14] (for $k := 0$), it follows that

Proposition 1 ([14]). Let $\mathcal{T} := \text{E-G}_\infty\text{A}^\omega + \text{QF-AC} + \text{WKL}$ and $\forall f \exists n A_{qf}(f, n)$ a sentence as above. Furthermore, let φ a closed term of \mathcal{T} (of suitable type). Then the following rule holds:

$$\left\{ \begin{array}{l} \mathcal{T} \vdash \forall f \in \mathbb{N}^{\mathbb{N}} (\Pi_1^0\text{-CA}(\varphi(f)) \rightarrow \exists n \in \mathbb{N} A_{qf}(f, n)) \\ \Rightarrow \text{there exists a primitive recursive (in the sense Kleene) functional } \Phi \text{ s.t.} \\ \widehat{\text{E-PA}}^\omega \upharpoonright \vdash \forall f \in \mathbb{N}^{\mathbb{N}} A_{qf}(f, \Phi(f)). \end{array} \right.$$

In this rule, we may add an arbitrary set of true purely universal sentences \mathcal{P} as additional axioms to both \mathcal{T} and $\widehat{\text{E-PA}}^\omega \upharpoonright$.

The main technical result in this paper establishes that over \mathcal{T} one can prove $\text{RT}_2^2(c)$ (i.e. Ramsey's theorem for pairs and a 2-coloring c) from a suitable instance $\Pi_1^0\text{-CA}(\tilde{\varphi}(c))$ of $\Pi_1^0\text{-CA}$, i.e.

Theorem 2 (see Theorem 21 below).

$$\mathcal{T} \vdash \forall c : [\mathbb{N}]^2 \rightarrow \mathbf{2} (\Pi_1^0\text{-CA}(\tilde{\varphi}(c)) \rightarrow \text{RT}_2^2(c)).$$

Instead of RT_2^2 we may have also RT_n^2 for any **fixed** number $n \geq 2$ of colors, where then $c : [\mathbb{N}]^2 \rightarrow \mathbf{n}$.

Here $[\mathbb{N}]^2$ denotes the set of unordered pairs in \mathbb{N} and \mathbf{n} the set $\{0, \dots, n-1\}$.

Combined with the previous result (and the fact that finitely many and even sequences of instances of $\Pi_1^0\text{-CA}$ can be encoded into a single instance) we obtain

Theorem 3 (see Theorem 23 below). Let φ, ψ be closed terms of \mathcal{T} (of suitable type). Then the following rule holds:

$$\left\{ \begin{array}{l} \mathcal{T} \vdash \forall f \in \mathbb{N}^{\mathbb{N}} (\Pi_1^0\text{-CA}(\varphi(f)) \wedge \forall k \in \mathbb{N} (\text{RT}_2^2(\psi(f, k))) \rightarrow \exists n \in \mathbb{N} A_{qf}(f, n)) \\ \Rightarrow \text{there exists a primitive recursive (in the sense Kleene) functional } \Phi \text{ s.t.} \\ \widehat{\text{E-PA}}^\omega \upharpoonright \vdash \forall f \in \mathbb{N}^{\mathbb{N}} A_{qf}(f, \Phi(f)). \end{array} \right.$$

Instead of RT_2^2 we, again, may have RT_n^2 for any fixed number n of colors.

We, furthermore, may add arbitrary true universal sentences as axioms to the theories in question.

Note that we cannot replace \mathcal{T} by $\widehat{\text{E-PA}}^\omega \upharpoonright$ or any other system containing either Σ_1^0 -induction (with function parameters) or the functional Φ_{it} as in such a system even Proposition 1 would be wrong, see [13].

For Π_2^0 -sentences $\forall m \in \mathbb{N} \exists n \in \mathbb{N} A_{qf}(m, n)$ one gets with Theorem 3 – using the well-known fact that $\widehat{\text{E-PA}}^\omega \upharpoonright$ is Π_2^0 -conservative over primitive recursive arithmetic (with quantifiers) PRA – as conclusion

$$\text{PRA} \vdash \forall m \in \mathbb{N} A_{qf}(m, \varphi(m)).$$

Let (for fixed n) RT_n^{2-} and $\Pi_1^0\text{-CA}^-$ be (the universal closures) of all instances $\text{RT}_n^2(s)$ and $\Pi_1^0\text{-CA}(t)$ for terms s, t containing only number parameters. Then we get

Corollary 4.

$$\mathcal{T} + \Pi_1^0\text{-CA}^- + \text{RT}_n^{2-}$$

is Π_2^0 -conservative over PRA.

Combined with further results from [14] it also follows $\mathcal{T} + \Pi_1^0\text{-CA}^- + \text{RT}_n^{2-}$ is Π_3^0 -conservative over $\text{PRA} + \Sigma_1^0\text{-IA}$.

Officially every variable in our system has a type (e.g. 0 for a natural number and 1 for a function $\mathbb{N} \rightarrow \mathbb{N}$, for details see [15]), but for simplicity of notation in the following we will denote by b, c, f, g, h, q number-theoretic functions of suitable arity and by $x, y, z, k, l, m, n, u, v$ natural numbers.

At a first look, it seems that the framework provided by \mathcal{T} is very restricted as only quantifier-free induction QF-IA (with parameters of arbitrary types) is included. However, from $\Pi_1^0\text{-CA}(\varphi(f))$ (for suitable φ) combined with QF-IA one obtains fixed sequences of instances

$$\Sigma_1^0\text{-IA}(f) := \left\{ \begin{array}{l} \forall l (\exists y (f(0, y, l) = 0) \wedge \forall x (\exists y (f(x, y, l) = 0) \rightarrow \exists y (f(x+1, y, l) = 0)) \\ \rightarrow \forall x \exists y (f(x, y, l) = 0)) \end{array} \right.$$

of Σ_1^0 -induction. Hence the theorem above also holds with

$$\Pi_1^0\text{-CA}(\varphi(f)) \wedge \forall k \in \mathbb{N} (\text{RT}_2^2(\psi(f, k)))$$

being replaced by

$$\Pi_1^0\text{-CA}(\varphi(f)) \wedge \forall k \in \mathbb{N} (\text{RT}_2^2(\psi(f, k))) \wedge \Sigma_1^0\text{-IA}(\chi(f)).$$

So, in particular, any instance of the schema of $\Sigma_1^0\text{-IA}$ restricted to Σ_1^0 -formulas that only has free number variables (short $\Sigma_1^0\text{-IA}^-$) is allowed. What is not permitted, however, is that the **result** of the application of $\Pi_1^0\text{-CA}(\varphi(f))$ (i.e. the

comprehension function) or of $\text{RT}_2^2(\psi(f, k))$ (i.e. the monochromatic set given by its characteristic function) is used in forming a Σ_1^0 -instance of induction.

Combined with $\text{QF-AC}^{\mathbb{N}, \mathbb{N}}$ one can use $\Pi_1^0\text{-CA}(f)$ even to obtain every instance of Δ_2^0 -comprehension as well as Π_1^0 -countable choice for numbers and hence every fixed sequence of instances of Δ_2^0 -induction and Π_1^0 -bounded collection, where the latter is defined as

$$\begin{aligned} \Pi_1^0\text{-CP}(f) &:= \\ \forall k, l \left(\forall x < l \exists y \forall z (f(k, x, y, z) = 0) \rightarrow \exists y^* \forall x < l \exists y < y^* \forall z (f(k, x, y, z) = 0) \right) \end{aligned}$$

(see [13]).

Here $\text{QF-AC}^{\mathbb{N}, \mathbb{N}}$ denotes the special case of QF-AC where both variables (n, f) are natural numbers.

Finally, we note that relative to \mathcal{T} fixed sequences of instances of the Bolzano-Weierstraß principle and even the Ascoli lemma can be proven from $\Pi_1^0\text{-CA}(\xi)$ for a suitable ξ (see [11]).

What all this indicates is that from the perspective of unwinding the computational content of concrete proofs based on RT_2^2 (and even RT_n^2 for fixed n) the computational complexity of that content will in most practical cases not go beyond primitive recursive complexity.

Theorem 2 is established by a careful analysis of the proof of Ramsey's theorem for pairs due to Erdős and Rado [4] which first yields that relative to suitable (sequences of) instances of Π_1^0 -induction (or – equivalently – Σ_1^0 -induction) with the coloring c as the only free function variable (so that these instances can be covered as discussed above)

$$\Sigma_1^0\text{-WKL}(\varphi(c)) \rightarrow \text{RT}_2^2(c)$$

for a suitable elementary functional φ . Here $\Sigma_1^0\text{-WKL}$ is König's lemma for 0/1-trees which are given by a Σ_1^0 -formula. $\Sigma_1^0\text{-WKL}(\varphi(c))$ (as well as the inductions needed) is then reduced using $\Pi_1^0\text{-CA}(\tilde{\varphi}(c))$ (for a suitable functional $\tilde{\varphi}$) to WKL and quantifier-free induction which both are available in \mathcal{T} .

2. ELIMINATION OF MONOTONE SKOLEM FUNCTIONS

In [13, 14] the second author developed a technique for the elimination of monotone Skolem functions that allows one to calibrate the arithmetical strength of fixed (sequences of) instances of various comprehension and choice principles over systems such as $\text{E-G}_\infty\text{A}^\omega$. In this section we collect the results of this type that will be used later.

The next result immediately follows (as special case for $k := 1$) from the proofs of Corollaries 4.4 and 4.5 in [14]:

Proposition 5 ([14]). Let $A_{qf}(f, g, n) \in \mathcal{L}(\text{E-G}_\infty\text{A}^\omega)$ be a quantifier-free formula which contains only the function variables f, g and the number variable n free.

Furthermore, let φ, ψ be functionals (of suitable type) that are definable in $E\text{-G}_\infty A^\omega$. Then the following rule holds

$$\left\{ \begin{array}{l} E\text{-G}_\infty A^\omega + \text{QF-AC} + \text{WKL} \vdash \forall f \forall g \leq \varphi(f) (\Pi_1^0\text{-CA}(\psi(f, g)) \rightarrow \exists n A_{qf}(f, g, n)) \\ \text{then one can extract a closed term } \Phi \text{ of } \widehat{E\text{-PA}}^\omega \upharpoonright \text{ such that} \\ \widehat{E\text{-PA}}^\omega \upharpoonright \vdash \forall f \forall g \leq \varphi(f) \exists n \leq \Phi(f) A_{qf}(f, g, n). \end{array} \right.$$

Here ‘ \leq ’ for functions is defined pointwise.

Proof: As in the proof of Corollary 4.5 in [14], we can replace WKL by the principle F^- and then use elimination of extentionality (see e.g. [15], the restrictions on the types in QF-AC are made precisely to allow for this) to obtain

$$(G_\infty A^\omega + \text{QF-AC}) \oplus F^- \vdash \forall f \forall g \leq \varphi(f) (\Pi_1^0\text{-CA}(\psi(f, g)) \rightarrow \exists n A_{qf}(f, g, n)).$$

Then apply Corollary 4.4 (for $\Delta := \emptyset$) and note that for $k := 1$ the conclusion can be verified in (even the weakly extensional and intuitionistic version of) $\widehat{E\text{-PA}}^\omega \upharpoonright$. \square

Remark 6. The instance of Π_1^0 -comprehension in Proposition 5 may also depend on the results of instances of WKL:

WLK($\tau(f)$) is implied by $\exists b \leq 1 \forall x (\tilde{\tau}(f)(\bar{b}x) = 0)$ for a suitable term $\tilde{\tau}$ in $E\text{-G}_\infty A^\omega$, with

$$E\text{-G}_\infty A^\omega \vdash \forall f, x^* \exists b \leq 1 \forall x \leq x^* (\tilde{\tau}(f)(\bar{b}x) = 0),$$

see [15, Proposition 9.18] (note that the g in the proof of this proposition is definable in $E\text{-G}_\infty A^\omega$). Suppose now that $E\text{-G}_\infty A^\omega + \text{QF-AC} + \text{WKL}$ proves

$$\forall f \forall g \leq \varphi(f) \forall b \leq 1 (\forall x \tilde{\tau}(f)(\bar{b}x) = 0 \rightarrow (\Pi_1^0\text{-CA}(\xi(f, b)) \rightarrow \exists n A_{qf}(f, n))),$$

which is equivalent to

$$\forall f \forall g \leq \varphi(f) \forall b \leq 1 (\Pi_1^0\text{-CA}(\xi(f, b)) \rightarrow \exists n, x (\tilde{\tau}(f)(\bar{b}x) = 0 \rightarrow A_{qf}(f, g, n))).$$

Applying Proposition 5 yields bounds $x^* := \chi(f)$ and $n^* := \Phi(f)$ on x and n depending only on f , i.e.

$$\widehat{E\text{-PA}}^\omega \upharpoonright \vdash \forall f \forall g \leq \varphi(f) (\exists b \leq 1 \forall x \leq \chi(f) (\tilde{\tau}(f)(\bar{b}x) = 0) \rightarrow \exists n \leq \Phi(f) A_{qf}(f, g, n))$$

and so, finally,

$$\widehat{E\text{-PA}}^\omega \upharpoonright \vdash \forall f \forall g \leq \varphi(f) \exists n \leq \Phi(f) A_{qf}(f, g, n).$$

Instead of fixed instances of $\Pi_1^0\text{-CA}$ also fixed sequences of such instances, i.e. fixed instances of

$$\Pi_1^0\text{-CA}^*(f) := \forall l \exists g \forall x (g(x) = 0 \leftrightarrow \forall y (f(l, x, y) = 0))$$

are covered since (provably in $E\text{-G}_\infty A^\omega$)

$$\Pi_1^0\text{-CA}(\varphi(f)) \rightarrow \Pi_1^0\text{-CA}^*(f),$$

where $\varphi(f) := f(j_1 x, j_2 x, y)$ for some unpairing functions j_1, j_2 .

We now consider sequences of Π_1^0 -instances of countable choice for numbers:

$$\Pi_1^0\text{-AC}(f) \equiv \forall l (\forall x \exists y \forall z (f(l, x, y, z) = 0) \rightarrow \exists g \forall x, z (f(l, x, g(x), z) = 0)).$$

$\Pi_1^0\text{-AC}(f)$ can be reduced to $\Pi_1^0\text{-CA}(g)$ uniformly by

Proposition 7 ([13]).

$$\text{E-G}_\infty\text{A}^\omega + \text{QF-AC}^{\mathbb{N}, \mathbb{N}} \vdash \forall f (\Pi_1^0\text{-CA}(\varphi(f)) \rightarrow \Pi_1^0\text{-AC}(f))$$

for a suitable elementary functional φ .

Similarly, one has:

Proposition 8 ([14]).

$$\text{E-G}_\infty\text{A}^\omega + \text{QF-AC}^{\mathbb{N}, \mathbb{N}} \vdash \forall f, g (\Pi_1^0\text{-CA}(\varphi(f, g)) \rightarrow \Delta_2^0\text{-CA}(f, g))$$

for a suitable φ , where

$$\Delta_2^0\text{-CA}(f, g) \equiv \begin{cases} \forall l (\forall x ([\forall u \exists v (f(l, x, u, v) = 0) \\ \leftrightarrow \exists m \forall n (g(l, x, m, n) = 0)]) \\ \rightarrow \exists h \forall x (h(x) = 0 \leftrightarrow \forall u \exists v (f(l, x, u, v) = 0))). \end{cases}$$

As a consequence of propositions 7 and 8 we obtain

Proposition 9 ([14]). Proposition 5 also holds with $\Pi_1^0\text{-AC}(\chi(f, g))$ and $\Delta_2^0\text{-CA}(\zeta_1(f, g), \zeta_2(f, g))$ in addition to $\Pi_1^0\text{-CA}(\psi(f, g))$ (and likewise for sequences of instances of $\Delta_2^0\text{-IA}$ and $\Pi_1^0\text{-CP}$).

With the restriction P^- of second order principles P to instances with at most number parameters as discussed in the introduction we can formulate the next proposition which follows (as special case for $k := 1$) from Corollaries 4.8 and 4.10 in [14]:

Proposition 10 ([14]). $\text{E-G}_\infty\text{A}^\omega + \text{QF-AC} + \text{WKL} + \Delta_2^0\text{-CA}^- + \Pi_1^0\text{-AC}^-$ is Π_3^0 -conservative over $\text{PRA} + \Sigma_1^0\text{-IA}$ and Π_4^0 -conservative over $\text{PRA} + \Pi_1^0\text{-CP}$.

3. TREES AND KÖNIG'S LEMMA

Definition 11 (Tree).

- (1) A partial order on the natural numbers \prec is called *tree* if for every $x \in \mathbb{N}$ the set of all predecessors $pd(x) := \{y \in \mathbb{N} \mid y \prec x\}$ is well-ordered.
- (2) A maximal linear order in \prec is called *branch*.
- (3) A tree \prec is called *finitely branching* if for all $x \in \mathbb{N}$ the set of all immediate successors $succ(x) := \{y \in \mathbb{N}, x \prec y \wedge (\nexists z (x \prec z \wedge z \prec y))\}$ is finite.
A tree is called *n-branching* if $|succ(x)| \leq n$ for all $x \in \mathbb{N}$.

Definition 12 (König's Lemma). König's Lemma is the statement that every infinite, finitely branching tree contains an infinite branch.

3.1. Fragments of König's Lemma and Formalizations. We formalize trees as characteristic functions of finite, initial segments of branches in a tree, i.e. a tree \prec is described by f if

$$\begin{aligned} f(\langle \rangle) &= 0 \\ f(\langle x \rangle) &= 0 \quad \text{iff} \quad x \text{ is } \prec\text{-minimal} \\ f(\langle n_1, \dots, n_k, x \rangle) &= 0 \quad \text{iff} \quad f(\langle n_1, \dots, n_k \rangle) = 0 \text{ and } x \in \text{succ}_{\prec}(n_k). \end{aligned}$$

We define $*$, $\langle \rangle$, \bar{b} using a suitable surjective sequence coding, for details see [15].

Definition 13 (Weak König's Lemma $\text{WKL}(\varphi)$).

$$\text{WKL}(\varphi): T(\varphi) \wedge \forall x^0 \exists s^0 (lth(s) = x \wedge \varphi(s)) \rightarrow \exists b \leq \lambda z.1 \forall x \varphi(\bar{b}x),$$

where T asserts that φ describes a 0, 1-tree with respect to the prefix relation \sqsubseteq

$$T(\varphi) := \forall s, r (\varphi(s * r) \rightarrow \varphi(s)) \wedge \forall s, x (\varphi(s * \langle x \rangle) \rightarrow x \leq 1).$$

Definition 14 (Bounded König's Lemma ($\text{WKL}^*(\varphi, h)$)).

$$\text{WKL}^*(\varphi, h): T^*(\varphi, h) \wedge \forall x^0 \exists s^0 (lth(s) = x \wedge \varphi(s)) \rightarrow \exists b \leq h \forall x \varphi(\bar{b}x),$$

where T^* asserts that φ describes a tree bounded by h

$$T^*(\varphi, h) := \forall s, r (\varphi(s * r) \rightarrow \varphi(s)) \wedge \forall s, x (\varphi(s * \langle x \rangle) \rightarrow x \leq h(lth(s))).$$

We denote by $\Sigma_1^0\text{-WKL}(f)$ resp. $\Sigma_1^0\text{-WKL}^*(f)$ weak/bounded König's Lemma with $\varphi(s) \equiv \exists z f(z, s) = 0$ and $\Sigma_1^0\text{-WKL}^{(*)} := \forall f \Sigma_1^0\text{-WKL}^{(*)}(f)$.

Proposition 15. In $\text{E-G}_\infty\text{A}^\omega$ every instance of bounded König's Lemma is equivalent to an instance of weak König's Lemma (WKL). Moreover, every instance of $\Sigma_1^0\text{-WKL}^*$ can be proven from an instance of $\Sigma_1^0\text{-WKL}$.

Proof. Simpson proves this equivalence in the system RCA_0 in [19, IV.1.3]. This proof can be carried out in $\text{E-G}_\infty\text{A}^\omega$. For $\varphi \in \Sigma_1^0$, this property is preserved. \square

Remark 16.

$$\text{E-G}_\infty\text{A}^\omega + \text{QF-AC}^{\mathbb{N}, \mathbb{N}} \vdash \Sigma_1^0\text{-WKL}(\xi(f)) \rightarrow \Pi_1^0\text{-CA}(f),$$

since $\Sigma_1^0\text{-WKL}(\sigma(g))$ implies $\Pi_2^0\text{-WKL}(g)$ for a suitable term σ , see [12, Proposition 3.3] or [19, proof of Lemma IV.4.4] and note that $\text{E-G}_\infty\text{A}^\omega + \text{QF-AC}^{\mathbb{N}, \mathbb{N}} \vdash \Sigma_1^0\text{-CP}$. $\Pi_2^0\text{-WKL}(\tau f)$ implies $\Pi_1^0\text{-CA}(f)$, see [21, §5].

Combined with the discussion at the end of section 1 it follows that over $\text{E-G}_\infty\text{A}^\omega + \text{QF-AC}^{\mathbb{N}, \mathbb{N}} + \text{WKL}$ each instance of $\Sigma_1^0\text{-WKL}$ is equivalent to an instance of $\Pi_1^0\text{-CA}$ and vice versa.

4. RAMSEY'S THEOREM

Now we turn to Ramsey's Theorem for pairs. In this section we will present two proofs of it. The first proof is the original one due to Ramsey [17], the second is due to Erdős and Rado [4, 10.2].

Definition 17.

- (1) $[X]^k := \{Y \subseteq X \mid |Y| = k\}$.
- (2) An n -coloring c of $[X]^k$ is a map $c: [X]^k \rightarrow \mathbf{n}$.
- (3) A set $H \subseteq X$ is called *monochromatic* under c if c is constant on $[H]^k$.
- (4) Let (X, \prec) be a partial order and c an n -coloring of $[X]^2$. A set $H \subseteq X$ is called *min-monochromatic* under c if for all $i \in H$ the map $c_i(x) := c(\{i, x\})$ is constant on $\{x \in H: i \prec x\}$.

Definition 18 (Ramsey's Theorem). For all k, n and every n -coloring c of $[\mathbb{N}]^k$ exists an infinite set $H \subseteq \mathbb{N}$, such that H is monochromatic under c .

RT_n^k denotes Ramsey's Theorem for n -colorings of $[\mathbb{N}]^k$ and, $\text{RT}_{<\infty}^k$ is defined as $\forall n \text{RT}_n^k$.

The proofs we are going to present share the same structure: First an infinite min-monochromatic set is constructed, then using RT_n^1 one finds an infinite monochromatic set.

Ramsey's proof is simpler and seemingly elementary, but it cannot even be formalized in ACA_0 , see [19, p. 123]. Therefore this proof is unusable for a detailed analysis of the proof-theoretic strength of RT_n^2 .

Erdős' and Rado's proof can be formalized in ACA_0 (see [19, Lemma III.7.4]). It uses König's Lemma, which is open for a detailed analysis in this case.

Ramsey's Proof. Fix an n -coloring $c: [\mathbb{N}]^2 \rightarrow \mathbf{n}$.

We construct an enumeration $(x_j)_{j \in \mathbb{N}}$ of an infinite min-monochromatic set.

Define $c_y(x) := c(\{y, x\})$.

- Set $x_0 := 0$.
- Using RT_n^1 we find an infinite set $X_1 \subseteq \mathbb{N} \setminus \{x_0\}$, such that X_1 is monochromatic under c_0 . Set $x_1 := \min X_1$.
- Similarly we find an infinite set $X_2 \subseteq X_1 \setminus \{x_1\}$, such that X_2 is monochromatic under c_{x_1} . Set $x_2 := \min X_2$.
- ...

Iterating this process gives a sequence $(x_j)_{j \in \mathbb{N}}$. By construction $X := \{x_0, x_1, \dots\}$ is min-monochromatic under c .

Define $c' : X \rightarrow \mathbf{n}$ with $c'(x_j) := c(\{x_j, x_{j+1}\})$. c' is well-defined since the sequence $(x_j)_j$ is injective.

Using RT_n^1 we find an infinite $H \subseteq X$ such that H is monochromatic under c' . Since H is min-monochromatic under c we get for all $x, y \in H, x < y$

$$c(\{x, y\}) = c'(x) = c'(H).$$

In other words, H is monochromatic under c . □

Erdős' and Rado's Proof. The notation of this proofs follows [5].

Fix an n -coloring $c : [\mathbb{N}]^2 \rightarrow \mathbf{n}$.

Let $c_k : \mathbf{k} \rightarrow \mathbf{n}$ be defined as $x \mapsto c(\{x, k\})$.

Now define recursively a partial order \prec on \mathbb{N} :

- $0 \prec 1$
- If \prec is already defined on \mathbf{m} , then let

$$P_k := \{x \in \mathbf{m} \mid x \prec k\} \quad \text{for } k \in \mathbf{m}.$$

Now, to extend \prec to $\mathbf{m} + \mathbf{1}$, for $k \in \mathbf{m}$ set

$$k \prec m \quad \text{iff} \quad c_k|_{P_k} = c_m|_{P_k}.$$

Claim:

- (i) $\prec \subseteq <_{\mathbb{N}}$, in particular $P_k = pd(k)$.
- (ii) $0 \prec x$ for all $x \in \mathbb{N} \setminus \{0\}$.
- (iii) \prec is transitive.
- (iv) On $pd(m)$ the relations $<_{\mathbb{N}}$ and \prec describe the same order, i.e. for $x, y \in pd(m)$

$$x < y \quad \text{iff} \quad x \prec y.$$

(i), (ii) follow immediately from the definition of \prec .

(iii):

We prove the statement $(x \prec y \text{ and } y \prec z) \Rightarrow x \prec z$ by induction on z .

The base case $z = 0$ is trivial because of (i).

Assume that transitivity holds for all $z' < z$. Then

$$\begin{aligned} x \prec y \text{ and } y \prec z &\Rightarrow c_x|_{P_x} = c_y|_{P_x}, c_y|_{P_y} = c_z|_{P_y} \\ &\text{and } P_x \subseteq P_y \quad (\text{induction hypothesis for } y < z) \\ &\Rightarrow c_x|_{P_x} = c_y|_{P_x} = c_z|_{P_x} \\ &\Rightarrow x \prec z. \end{aligned}$$

(iv):

“ \Leftarrow ”: follows from (i).

“ \Rightarrow ”: By (ii) the case $x = 0$ is trivial. Let $x \neq 0$.

Proof by induction on m :

- $m = 0$ is obvious.
- Let $m > 0$, $x, y \in pd(m)$ with $x < y$ and assume the statement holds for all $m' < m$.

Let i be the $<$ -maximal natural number such that $i \prec x$ and $i \prec y$ (such an i exists because of $0 \prec x, y$ by (ii)). Let p be an immediate \prec -successor of i comparable with m (such a p exists because of $i \prec x \prec m$). From $i \prec y \prec m$ and $i \prec p \prec m$ we get

$$c_y(i) = c_m(i) = c_p(i).$$

Using the induction hypothesis for $m' = p$, we deduce that all $i' \prec p$ are comparable with i , in particular $p \in \text{succ}(i)$ and

$$P_p = P_i \cup \{i\}.$$

Since $i \prec y$ and $c_y(i) = c_p(i)$, this shows $p \prec y$ (the case $p = y$ is impossible). Analogously, it follows that $p \prec x$ or $p = x$. The maximality of i renders the case $p \prec x$ impossible, so $p = x$ and in particular $x \prec y$.

By (iv) the relation \prec defines a tree on \mathbb{N} .

By definition, every branch of \prec is min-monochromatic under c .

The tree is n -branching (in particular finitely branching) since for all $x, y \in \text{succ}(i)$ such that $x < y$ the induced colorings c_x and c_y must differ at i . Otherwise $x \prec y$ since $c_x|_{P_i} = c_y|_{P_i}$ and $P_x = P_i \cup \{i\}$.

By König's Lemma we find an infinite min-monochromatic branch B . As in Ramsey's proof we construct using RT_n^1 an infinite monochromatic set H under c . \square

Note that we cannot simply reduce the application of König's Lemma in this proof (or in Simpson's proof [19, Lemma III.4.7]) to WKL using an instance of $\Pi_1^0\text{-AC}$ since we need $\Sigma_1^0\text{-IA}$ depending on the result of a $\Pi_1^0\text{-AC}$ application to prove that the tree is bounded. A bounding function for the tree can be constructed using only fixed instances of $\Sigma_1^0\text{-IA}$, $\Pi_1^0\text{-IA}$ and $\Pi_1^0\text{-AC}$, but this construction depends crucially on the special structure of the Erdős-Rado-tree (see [16]). However, we will follow here a slightly different approach.

4.1. Formalized proof of RT_n^2 . In the following we formalize the proof of Erdős and Rado and show Theorem 2. We proof this theorem for every fixed number $n \geq 2$ of colors, since the usual equivalence between RT_2^2 and RT_n^2 does not hold for individual instances.

In $\text{E-G}_\infty A^\omega$ we represent an n -coloring $c: [\mathbb{N}]^2 \rightarrow \mathbf{n}$ using a mapping $\hat{c}: \mathbb{N} \times \mathbb{N} \rightarrow \mathbf{n}$ such that $\hat{c}(x, y) = \hat{c}(y, x) = c(\{x, y\})$. We formalize RT_n^2 as follows

$$\begin{aligned} (\text{RT}_n^2): \forall c: \mathbb{N} \times \mathbb{N} \rightarrow \mathbf{n} \exists f \leq \lambda x.1 \exists i < n (\forall k \exists x > k f(x) = 0 \\ \wedge \forall x, y (x \neq y \wedge f(x) = 0 \wedge f(y) = 0 \rightarrow \hat{c}(x, y) = i)), \end{aligned}$$

$$\text{where } \hat{c}(x, y) = \begin{cases} c(x, y) & x \leq y, \\ c(y, x) & x > y. \end{cases}$$

RT_n^2 expresses that f is the characteristic functions of an infinite set in which every (unordered) pair $\{x, y\}$ is mapped to the color i .

$\text{RT}_n^2(t)$ denotes RT_n^2 for a fixed coloring t , RT_n^{2-} denotes the set of all instances of $\text{RT}_n^2(t)$, where the only free variables of t are of degree 0, i.e. number variables. We omit n , when no confusion can arise.

Lemma 19. For every coloring $c: \mathbb{N} \times \mathbb{N} \rightarrow \mathbf{n}$ the partial order \prec as in the proof of Erdős and Rado can be defined in $\text{E-G}_\infty A^\omega$.

$E\text{-G}_\infty A^\omega$ proves that \prec -chains are min-monochromatic and the properties (i)–(iv), i.e.

- (i) $\forall x, y (x \prec y \rightarrow x < y)$,
- (ii) $\forall x > 0 (0 \prec x)$,
- (iii) $\forall x, y, z (x \prec y \wedge y \prec z \rightarrow x \prec z)$,
- (iv) $\forall m, x, y (y \prec m \rightarrow (x \prec y \leftrightarrow x \prec m \wedge x < y))$.

Proof. We may assume $c(x, y) = c(y, x)$.

Define

$$\begin{aligned}\tilde{q}(0) &:= \langle \rangle, \\ \tilde{q}(1) &:= \langle 0 \rangle, \\ \tilde{q}(m+1) &:= \langle q_0^{m+1}, \dots, q_m^{m+1} \rangle,\end{aligned}$$

$$\text{where } q_k^{m+1} := \begin{cases} 0, & \text{if } \forall x < k ((\tilde{q}(k))_x = 0 \rightarrow c(k, x) = c(m+1, x)), \\ 1, & \text{otherwise} \end{cases}$$

for $k := 0, \dots, m$.

By definition $\tilde{q}(m) \leq \overline{\lambda x. \mathbb{1}(m)}$.

The mapping

$$q(x, y) := \begin{cases} \tilde{q}(y)_x & x < y \\ 1 & x \geq y \end{cases}$$

is the characteristic function of \prec . Hence the relation \prec can be defined with elementary recursion and so, in particular, in $E\text{-G}_\infty A^\omega$. Set

$$x \prec y := q(x, y) = 0.$$

- (i), (ii) immediately follow from definition of \prec resp. the mapping q .
- (iii) is (using (i)) equivalent to

$$\forall z \forall y < z, x < y (x \prec y \wedge y \prec z \rightarrow x \prec z).$$

We prove this statement using quantifier-free course-of-value induction on z .

The base case is trivial.

Assume that the statement holds for $z' < z$.

$$\begin{aligned}x \prec y \wedge y \prec z \\ \rightarrow [(\forall i < x (i \prec x \rightarrow c(x, i) = c(y, i))) \wedge (\forall i < y (i \prec y \rightarrow c(y, i) = c(z, i)))]\end{aligned}$$

using induction hypothesis for $y < z$

$$\begin{aligned}\rightarrow [(\forall i < x (i \prec x \rightarrow c(x, i) = c(y, i))) \wedge (\forall i < y (i \prec x \rightarrow c(y, i) = c(z, i)))] \\ \rightarrow [\forall i < x (i \prec x \rightarrow c(x, i) = c(z, i))] \\ \rightarrow x \prec z.\end{aligned}$$

(iv):

The “ \rightarrow ”-direction follows from (i) and (iii).

The “ \leftarrow ”-direction is (using (i)) equivalent to

$$\forall m \forall x < m, y < m (x \prec m \wedge y \prec m \wedge x < y \rightarrow x \prec y).$$

We prove this statement using quantifier-free course-of-value induction on m .

The base step is trivial.

Assume that the statement holds for all $m' < m$:

For $x = 0$ the statement is obvious. Hence we assume $x \neq 0$. Let $x \prec m$, $y \prec m$ and $x < y$.

$$x \neq 0$$

$$\xrightarrow{(ii)} \exists i < x (i \prec x \wedge i \prec y) \quad (\text{e.g. } i = 0)$$

$$\xrightarrow{\mu_b} \exists i < x (i \prec x \wedge i \prec y \wedge \underbrace{\forall i' < x ((i' \prec x \wedge i' \prec y) \rightarrow i' \leq i)}_{\equiv: i \text{ maximal}})$$

$$\rightarrow \exists i < x (i \prec x \wedge i \prec y \wedge i \text{ maximal} \wedge \exists p < m (p \prec m \wedge i \prec p)) \quad (\text{e.g. } p = x)$$

$$\xrightarrow{\mu_b} \exists i < x (i \prec x \wedge i \prec y \wedge i \text{ maximal}$$

$$\wedge \exists p < m (\underbrace{p \prec m \wedge i \prec p \wedge \forall p' < m (p' \prec m \wedge i \prec p' \rightarrow p' \geq p)}_{p \text{ minimal with } i \prec p \prec m})).$$

Using (iii), we deduce $i \prec m$. Since $y \prec m$ and $i \prec y$, this gives $c(y, i) = c(m, i)$.

From $p \prec m$ and $i \prec p$ it follows $c(p, i) = c(m, i)$. Therefore

$$(1) \quad c(y, i) = c(p, i).$$

From the induction hypothesis (for p) and (i) we obtain

$$(2) \quad \forall j, j' ((j \prec p \wedge j' \prec p) \rightarrow (j \prec j' \leftrightarrow j < j')).$$

We claim that p is an immediate successor of i . In other words, no i' exists such that $i \prec i' \prec p$. Suppose such an i' exists. Then (iii) gives $i \prec i' \prec m$. As p is minimal with this property we get $i' \geq p$. This contradicts (together with (i)) the assumption $i' \prec p$.

Combining this with (2) we see

$$(3) \quad \forall i' (i' \prec p \rightarrow (i' = i \vee i' \prec i)).$$

Since $i \prec y, p$ we get $c(p, i') = c(i, i') = c(y, i')$ for all $i' \prec i$.

This, (1) and, (3) shows $c(p, i') = c(y, i')$ for all $i' \prec p$ and, in particular, $p \prec y$ ($p = y$ is impossible because of $p \leq x < y$).

This implies

$$\exists i < x (i \prec x \wedge i \prec y \wedge i \text{ maximal} \wedge \exists p < m (p \prec y \wedge p \in \text{succ}(i))).$$

Analogously, we deduce $p = x$. Here the maximality of i renders the case $p \prec x$ impossible. Put together, we obtain

$$\exists i < x (i \prec x \wedge i \prec y \wedge i \text{ maximal} \wedge \exists p < m (p \prec y \wedge p = x))$$

and so

$$x \prec y.$$

□

Lemma 20. For every fixed $n \geq 2$ there are closed terms ξ_1 and ξ_2 such that

$$\begin{aligned} \text{E-G}_\infty\text{A}^\omega + \text{QF-AC}^{\mathbb{N},\mathbb{N}} \vdash \forall c: \mathbb{N} \times \mathbb{N} \rightarrow \mathbf{n} \left(\Pi_1^0\text{-CA}(\xi_1 c) \wedge \Sigma_1^0\text{-WKL}(\xi_2 c) \right. \\ \left. \rightarrow \exists b (b_0 = 0 \wedge \forall i b(i+1) \in \text{succ}(bi)) \right). \end{aligned}$$

Proof. Notation as in the proof of the preceding lemma.

Define:

$$\begin{aligned} h(0, q^0, c^0) &:= \langle \rangle = 0 \\ h(m+1, q^0, c^0) &:= h(m, q, c) * \begin{cases} \langle (c)_{m+1} \rangle & \text{if } (q)_{m+1} = 0 \\ \langle \rangle = 0 & \text{else} \end{cases} \\ h(k, q, c) &\leq c \\ g(m) &:= h(m, \Phi_{\langle \rangle}(\lambda x. q(x, m), m), \Phi_{\langle \rangle}(\lambda x. c(x, m), m)) \end{aligned}$$

The function h deletes the entries i in c , where $(q)_i \neq 0$ holds. Hence $g(m) = \langle c(m, i_0), \dots, c(m, i_k) \rangle$, where $i_0 \prec i_1 \prec \dots \prec i_k$ are the predecessors of m ordered by \prec . Note h and g can be defined in $\text{E-G}_\infty\text{A}^\omega$.

By definition of g

$$(4) \quad (g(x))_i < n,$$

$$(5) \quad x \prec y \rightarrow g(x) \sqsubset g(y).$$

We deduce:

$$\begin{aligned} g(z) &= m * \langle x \rangle \\ \xrightarrow{\mu_b} \exists v < z (g(z) = m * \langle x \rangle \wedge \underbrace{v < z \wedge \forall v' < z (v' \prec z \rightarrow v' \leq v)}_{v \text{ maximal with } v \prec z}) \\ &\rightarrow \exists v < z (g(z) = m * \langle x \rangle \wedge z \in \text{succ}(v)) \\ \xrightarrow{(iv)} \exists v < z (g(z) = m * \langle x \rangle \wedge z \in \text{succ}(v) \wedge \forall x < v (x \prec v \leftrightarrow x \prec z)) \\ &\rightarrow \exists v < z (g(z) = m * \langle x \rangle \wedge z \in \text{succ}(v) \wedge \tilde{q}(v) \sqsubset \tilde{q}(z)) \end{aligned}$$

since v is maximal with $v \prec z$, (i) yields $(\tilde{q}(z))_i = 0$ for all $i \in \{v+1, \dots, z-1\}$.

This gives us

$$\exists v < z (g(z) = m * \langle x \rangle \wedge z \in \text{succ}(v) \wedge g(v) = m)$$

and, in particular,

$$\exists v (g(v) = m \wedge z \in \text{succ}(v)).$$

We conclude

$$(6) \quad \forall n, x, z (g(z) = m * \langle x \rangle \rightarrow \exists v < z (g(v) = m \wedge z \in \text{succ}(v))).$$

We proceed to prove that g is injective by showing

$$(7) \quad \forall l \forall x, y (x \neq y \wedge lth(g(x)) = l \rightarrow g(x) \neq g(y))$$

using Π_1^0 -induction on l . Note that the induction formula can be written as $\Pi_1^0\text{-IA}(\xi_1 c)$ for a suitable ξ_1 .

The base case is an immediate consequence of (ii) and the definition of g .

Assume that (7) holds for l .

$$\begin{aligned} & \exists x, y (x \neq y \wedge lth(g(x)) = l + 1 \wedge g(x) = g(y)) \\ \xrightarrow{(6)} & \exists x, y \exists x', y' (x \neq y \wedge lth(g(x)) = l + 1 \wedge x \in succ(x') \wedge y \in succ(y') \wedge \\ & g(x) = g(y) \wedge g(x') \sqsubset g(x) \wedge lth(g(x')) = l \wedge g(y') \sqsubset g(y) \wedge lth(g(y')) = l) \\ \xrightarrow{\text{IH}} & \exists x, y \exists x' (x \neq y \wedge lth(g(x)) = l + 1 \wedge x, y \in succ(x') \wedge g(x) = g(y)) \end{aligned}$$

Since x and y are immediate successors of x' and $c(x, x') = (g(x))_l = (g(y))_l = c(y, x')$, either x, y are equal or comparable. The former case contradicts our assumption, the later together with (5) the fact that $g(x) = g(y)$. This finishes the proof of the injectivity of g .

The injectivity of g together with (6) yields

$$\forall z, v (g(z) = m * \langle x \rangle \wedge g(v) = m \rightarrow v \prec z).$$

Using Π_1^0 -induction and Lemma 19 (iii) we conclude

$$\forall l \forall x, y (lth(g(y)) = l \wedge g(x) \sqsubset g(y) \rightarrow x \prec y).$$

Since g is definable in terms of $\text{E-G}_\infty \text{A}^\omega$ and c , the induction formula can be written as $\Pi_1^0\text{-IA}(\xi_2 c)$ for a suitable term ξ_2 .

Together with (5) this gives us

$$(8) \quad x \prec y \leftrightarrow g(x) \sqsubset g(y).$$

Using (6) it is clear that

$$\xi_3(c, x, s) := g(x) = s$$

defines a Σ_1^0 -tree bounded by $\lambda z.n$. By definition the tree is the image of g . As g is an injection from the natural numbers the tree is infinite. Applying $\Sigma_1^0\text{-WKL}(\xi_3 c)$ yields a branch b' with

$$\begin{aligned} \forall i \exists x g(x) &= \bar{b}'i \\ \xrightarrow{\text{QF-AC}} \exists b \forall i g(bi) &= \bar{b}'i \end{aligned}$$

This and (8) establishes the lemma.

Note the two instances of Π_1^0 -induction can be proven from $\Pi_1^0\text{-CA}(\xi_1 c)$ for a suitable ξ_1 , see section 2. \square

Theorem 21. For each fixed $n \geq 2$ there exists a closed term ξ , such that

$$\text{E-G}_\infty \text{A}^\omega + \text{QF-AC}^{\mathbb{N}, \mathbb{N}} + \text{WKL} \vdash \forall c: \mathbb{N} \times \mathbb{N} \rightarrow \mathbf{n} (\Pi_1^0\text{-CA}(\xi c) \rightarrow \text{RT}_n^2(c)).$$

Proof. As

$$\text{E-G}_\infty\text{A}^\omega + \text{WKL} \vdash \Pi_1^0\text{-CA}(\xi c) \rightarrow (\Pi_1^0\text{-CA}(\xi_1 c) \wedge \Sigma_1^0\text{-WKL}(\xi_2 c))$$

for a suitable ξ , Lemma 20 gives us an infinite branch b of the Erdős-Rado tree \prec . By definition of \prec , $b(\mathbb{N})$ is min-monochromatic under c and

$$\forall x, y \ (x < y \leftrightarrow x \prec y).$$

Define $c'(x) := c(bx, b(x+1))$. Since $b(\mathbb{N})$ is min-monochromatic, we get

$$\forall x \forall y > x \ (c(bx, by) = c'x).$$

By RT_n^1 there exists a color i occurring infinitely often. The set $H := \{bx \mid c'x = i\}$ is infinite and monochromatic under c , so $b^*k := t_{\exists x \leq k \ bx=k \wedge c'k=i}[k]$ forms a solution of RT_n^2 . \square

Remark 22. Using the tuple coding from section 2 it is obvious that, for a suitable ξ , $\Pi_1^0\text{-CA}(\xi((c_k)_k))$ proves a sequence of instances $(\text{RT}_n^2(c_k))_{k \in \mathbb{N}}$.

Note that the number of colors in such a sequence of instances of RT_n^2 has to be bounded. For an unbounded number of colors we would need $\text{RT}_{<\infty}^1$ in the proof of Theorem 21. But as $\text{RT}_{<\infty}^1$ is equivalent to $\Pi_1^0\text{-CP}$ (see [8]) it is not provable in $\text{E-G}_\infty\text{A}^\omega$.

5. RESULTS

Using Theorem 21 we can extend the theorems of section 2 by adding RT_n^2 :

Theorem 23. Let $A_{qf}(f, g, k) \in \mathcal{L}(\text{E-G}_\infty\text{A}^\omega)$ be a quantifier-free formula which contains only the variables f, g, k free. Furthermore, let φ, ψ, χ be functionals (of suitable type) that are definable in $\text{E-G}_\infty\text{A}^\omega$. Then for every fixed $n \geq 2$ the following rule holds

$$\left\{ \begin{array}{l} \text{E-G}_\infty\text{A}^\omega + \text{QF-AC} + \text{WKL} \\ \vdash \forall f \forall g \leq \varphi(f) \ (\Pi_1^0\text{-CA}(\psi(f, g)) \wedge \forall l \text{RT}_n^2(\chi(f, g, l)) \rightarrow \exists k A_{qf}(f, g, k)) \\ \text{then one can extract a closed term } \Phi \text{ of } \widehat{\text{E-PA}}^\omega \text{ such that} \\ \widehat{\text{E-PA}}^\omega \upharpoonright \vdash \forall f \forall g \leq \varphi(f) \ \exists k \leq \Phi(f) \ A_{qf}(f, g, k). \end{array} \right.$$

Proof. Proposition 5, Theorem 21 and Remark 22. \square

Theorem 24. Let $\mathcal{S} := \text{E-G}_\infty\text{A}^\omega + \text{QF-AC} + \text{WKL} + \Delta_2^0\text{-CA}^- + \Pi_1^0\text{-AC}^-$ and n fixed. Then the following holds

- (i) $\mathcal{S} + \text{RT}_n^{2-}$ is Π_2^0 -conservative over PRA,
- (ii) $\mathcal{S} + \text{RT}_n^{2-}$ is Π_3^0 -conservative over PRA + $\Sigma_1^0\text{-IA}$,
- (iii) $\mathcal{S} + \text{RT}_n^{2-}$ is Π_4^0 -conservative over PRA + $\Pi_1^0\text{-CP}$.

Proof. (i) follows from Corollary 4 and Theorem 21.

(ii) and (iii) follow from Proposition 10 and Theorem 21. \square

The bound in (ii) is sharp. Avigad constructed in [1] a Σ_3^0 -sentence provable from $\Pi_1^0\text{-CP}^-$ and hence from $\Pi_1^0\text{-AC}^-$ that is not provable in $\text{PRA} + \Sigma_1^0\text{-IA}$.

These theorems cannot be extended to $\text{RT}_{<\infty}^2$:

Proposition 25.

$$\text{E-G}_\infty\text{A}^\omega + \text{QF-AC} + \text{WKL} + \Delta_2^0\text{-CA}^- + \Pi_1^0\text{-AC}^- \not\vdash \text{RT}_{<\infty}^{2^-}$$

Proof. A sequence of instances of $\text{RT}_{<\infty}^2$ with unbounded number of colors is sufficient to prove the totality of (a version of) the Ackermann function, see [8, 6.12].

All instances of $\text{RT}_{<\infty}^2$ in the proof of this theorem are of the form $\text{RT}_{<\infty}^{2^-}$. Since the diagonal of the Ackermann Function cannot be primitive recursively bounded, the theorem follows from Theorem 3, Proposition 7 and, 8. \square

Remark 26. Our formalization of the proof of RT_n^2 also can be used to analyze the complexity of RT_n^2 relative to the comprehension used (in our case $\Sigma_1^0\text{-WKL}$) like Bellin did in [2] using Ramsey's proof. The proof of Lemma 20 yields the concrete instance of the comprehension needed as an elementary functional in the coloring c (namely the term ξ_3 derived from the construction of the Erdős-Rado-tree). As we are not using Ramsey's proof in our case, a weaker instance of comprehension suffices. It should be noted, though, that the main concern in [2] is to derive a parametric version of Ramsey's theorem that displays the common structural features of the (proofs of the) (infinite) Ramsey theorem, the finite Ramsey theorem and the Paris-Harrington theorem.

REFERENCES

1. Jeremy Avigad, *Notes on Π_1^1 -conservativity, ω -submodels, and collection schema*, Tech. report, Carnegie Mellon Department of Philosophy, 2002.
2. Gianluigi Bellin, *Ramsey interpreted: a parametric version of Ramsey's theorem*, Logic and computation (Pittsburgh, PA, 1987), Contemp. Math., vol. 106, Amer. Math. Soc., Providence, RI, 1990, pp. 17–37. MR 1057813
3. Peter Cholak, Carl G. Jockusch, Jr., and Theodore A. Slaman, *On the strength of Ramsey's theorem for pairs*, J. Symb. Log. **66** (2001), no. 1, 1–55. MR 1825173
4. Paul Erdős, András Hajnal, Attila Máté, and Richard Rado, *Combinatorial set theory: partition relations for cardinals*, Studies in Logic and the Foundations of Mathematics, vol. 106, North-Holland Publishing Co., Amsterdam, 1984. MR 795592
5. Ilijas Farah, *Lecture notes: Set theory and its applications*, York University, unpublished, 2008.
6. Denis R. Hirschfeldt, Carl G. Jockusch, Jr., Bjørn Kjos-Hanssen, Steffen Lempp, and Theodore A. Slaman, *The strength of some combinatorial principles related to Ramsey's theorem for pairs*, Computational Prospects of Infinity, Part II: Presented Talks (Chitat Chong, Qi Feng, Theodore A. Slaman, W. Hugh Woodin, and Yue Yang, eds.), Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore, World Scientific, 2008, pp. 143–162.
7. Denis R. Hirschfeldt and Richard A. Shore, *Combinatorial principles weaker than Ramsey's theorem for pairs*, J. Symbolic Logic **72** (2007), no. 1, 171–206. MR 2298478
8. Jeffrey L. Hirst, *Combinatorics in subsystems of second order arithmetic*, Ph.D. thesis, Pennsylvania State University, 1987.

9. Carl G. Jockusch, Jr., *Ramsey's theorem and recursion theory*, J. Symbolic Logic **37** (1972), 268–280. MR 0376319
10. Ulrich Kohlenbach, *Mathematically strong subsystems of analysis with low rate of growth of provably recursive functionals*, Arch. Math. Logic **36** (1996), no. 1, 31–71. MR 1462200
11. ———, *Arithmetizing proofs in analysis*, Logic Colloquium '96 (San Sebastián), Lecture Notes Logic, vol. 12, Springer, Berlin, 1998, pp. 115–158. MR 1674949
12. ———, *The computational strength of extensions of weak König's lemma*, Research Series RS-98-41, BRICS, "Department of Computer Science, University of Aarhus", December 1998, 23 pp.
13. ———, *Elimination of Skolem functions for monotone formulas in analysis*, Arch. Math. Logic **37** (1998), 363–390. MR 1634279
14. ———, *On the arithmetical content of restricted forms of comprehension, choice and general uniform boundedness*, Annals of Pure and Applied Logic **95** (1998), 257–285. MR 1650651
15. ———, *Applied proof theory: Proof interpretations and their use in mathematics*, Springer Monographs in Mathematics, Springer Verlag, 2008. MR 2445721
16. Alexander Kreuzer, *Der Satz von Ramsey für Paare und beweisbar rekursive Funktionen*, Diploma Thesis, Technische Universität Darmstadt, 2009.
17. Frank P. Ramsey, *On a problem of formal logic*, Proc. London Math. Soc. **s2-30** (1930), no. 1, 264–286.
18. David Seetapun and Theodore A. Slaman, *On the strength of Ramsey's theorem*, Notre Dame J. Formal Logic **36** (1995), no. 4, 570–582, Special Issue: Models of arithmetic. MR 1368468
19. Stephen G. Simpson, *Subsystems of second order arithmetic*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1999. MR 1723993
20. Ernst Specker, *Ramsey's theorem does not hold in recursive set theory*, Logic Colloquium '69 (Proc. Summer School and Colloq., Manchester, 1969), North-Holland, Amsterdam, 1971, pp. 439–442. MR 0278941
21. Anne S. Troelstra, *Note on the fan theorem*, J. Symbolic Logic **39** (1974), 584–596. MR 0384494

FACHBEREICH MATHEMATIK, TECHNISCHE UNIVERSITÄT DARMSTADT, SCHLOSSGARTENSTRASSE 7,
64289 DARMSTADT, GERMANY