

SEMINAR ON CONTINUITY IN SEMILATTICES (SCS)

NAME(S) Mislove	DATE	M	D	Y
		8	18	76

TOPIC SCS Memo of Lawson dated 7-12-76

REFERENCE Ditto

In the above mentioned memo, Lawson defines the subset  $L(S)$  for a compact semilattice  $S$  to be those points of  $S$  where  $S$  has small semilattices. He shows that  $L(S)$  is closed under arbitrary sups, and that, for all  $s \in S$ , we have  $s \in L(S)$  iff  $s = \sup \downarrow s$ . The question is then raised as to whether  $L(S) \in \underline{CL}$ . The following example shows that such need not be the case:

Step 1. Let  $R \in \underline{CS}$  such that, for all subsemilattices  $S$  of  $R$ ,  $\text{int } S \neq \emptyset$  implies that  $0 \in S$  (such examples exist; see, e.g., Lawson, "Lattices with no interval homomorphisms" Pac. Jour. 32, 459-465). We let  $R' = \mathbb{N} \times R$  in the lexicographic order, where  $\mathbb{N}$  has its natural total order. Then, in the product topology,  $R'$  is a locally compact semilattice. Moreover, any compact subset of  $R'$  intersects at most finitely many of the sets  $\{n\} \times R$ . Hence, if  $R'' = R' \cup \{1\}$  is the one-point compactification of  $R'$ , it follows that the sets of the form  $\uparrow(n,0)$  form a neighborhood basis for the topology at 1. Thus, if we let 1 act as an identity for  $R''$ , we have that  $R''$  is a compact semilattice. It is readily verified that  $\mathcal{Q} = \{((n,0), (n+1,1)) : n \in \mathbb{N}\} \cup \Delta(R'')$  is a closed congruence on  $R''$ , so that  $T = R''/\mathcal{Q}$  is a compact semilattice with identity.

In "picturesque" language,  $T$  is a stack of countably many copies of  $R$  with an identity at the top. We wish to determine  $L(T)$ . Clearly  $0 \in L(T)$ , and since the sets of the form  $\uparrow(n,0)$  form a neighborhood basis at 1 in  $R''$ , the sets  $\uparrow[n,0]$  form a neighborhood basis at 1 in  $T$  (where  $[n,0]$  denotes the  $\mathcal{Q}$ -class of  $(n,0)$  in  $T$ ), and so we conclude that  $1 \in L(T)$  also. Now, let  $r \in R$ , and consider  $[n,r]$ . If  $U$  is any semilattice neighborhood of  $[n,r]$  in  $T$ , then  $U \cap (\{n\} \times R)$  is a semilattice neighborhood of  $[n,r]$  in the subsemilattice  $\{n\} \times R$ , which is isomorphic to  $R$ . Hence,  $[n,0] \in U$  by the defining property of  $R$ . Furthermore, if  $n > 0$ , then  $[n,0] = [n-1,1]$ , and the same argument as that just given for  $[n,r]$  shows that any semilattice neighborhood of  $[n,0]$  must also contain  $[n-1,0]$ . From these two facts we conclude that, for any  $n > 0$ , any semilattice neighborhood of  $[n,r]$  must also contain  $[0,0]$ , the zero of  $T$ . Thus,  $L(T) = \{0,1\}$ .


West Germany: TH Darmstadt (Gierz, Keimel)  
U. Tübingen (Mislove, Visit.)

England: U. Oxford (Scott)

USA: U. California, Riverside (Stralka)  
LSU Baton Rouge (Lawson)  
Tulane U., New Orleans (Hofmann, Mislove)  
U. Tennessee, Knoxville (Carruth, Crawley)

Step 2. The semilattice in which we are interested is a subsemilattice of  $T \times T$ . Let  $S = \bigcup_n \left( \{([n,0],[m,r]) : n \leq m \in \mathbb{N}\} \cup \{([m,r],[n,0]) : n \leq m \in \mathbb{N}\} \right) \cup \Delta(T)$   
 $= \bigcup_n \left( (p_1^{-1}([n,0]) \cap \uparrow([n,0],[n,0]) \cup (p_2^{-1}([n,0]) \cap \uparrow([n,0],[n,0])) \cup \Delta(T) \right)$

where  $p_i: T \times T \rightarrow T$  is the projection on the  $i^{\text{th}}$  factor. Then, from the second definition, it is clear that  $S$  is a countable union of closed subsemilattices of  $T \times T$ . Moreover, for each  $n \in \mathbb{N}$ , all but finitely many of these subsemilattices are contained in  $\uparrow([n,0],[n,0])$ , and these upper sets form a neighborhood basis at  $(1,1)$ . It then follows that  $S$  is a closed subset of  $T \times T$ . To see that  $S$  is a subsemilattice, we choose  $s, s' \in S$ , and we assume that  $s = ([n,0],[m,r])$  and  $s' = ([k,t],[j,0])$  with  $n \leq m$  and  $j \leq k$ . Either  $j \leq n$  or  $n \leq j$ , and we assume the former. Then,  $j \leq n \leq m$ , so that  $[m,r][j,0] = [j,0]$ . Also,  $j \leq k, n$  implies that  $[j,0] \leq [n,0][k,r]$ . Hence,  $ss' = ([i,0],[j,0])$  with  $j \leq \inf k, n = i$ , so that  $ss' \in S$  in this case. Similar arguments take care of the other possibilities, so that  $S$  is indeed a subsemilattice of  $T \times T$ . Hence,  $S \in \underline{CS}$ , and we want to determine  $L(S)$ . First we give a picture of  $S$ :



Now, clearly  $(0,0) \in L(S)$ , and since  $(1,1) = \sup_n ([n,0],[n,0])$  and  $([n,0],[n,0]) \ll_S (1,1)$  (this is not true in  $T \times T$ !) for each  $n \in \mathbb{N}$ , we have that  $(1,1) \in L(S)$  also. Now, if  $n \in \mathbb{N}$ , then it follows that  $([n,0],[n,0]) \ll_S ([n,0],1)$ , and so  $([n,0],[m,0]) \ll_S ([n,0],1)$  for each  $m > n$ . Hence  $([n,0],1) = \sup_S \downarrow ([n,0],1)$ , so that  $([n,0],1) \in L(S)$ , and a similar argument shows that  $(1,[n,0]) \in L(S)$  for each  $n \in \mathbb{N}$ . Finally, for each  $n \in \mathbb{N}$ ,  $p_1^{-1}([n,0]) \cap \uparrow([n,0],[n,0])$  is isomorphic to  $T$ , so that the only possible points of  $L(S)$  in this subsemilattice are  $([n,0],1)$  and  $([n,0],0)$ . We have already seen that  $([n,0],1) \in L(S)$ , and since  $([n,0],0) = ([n,0],[n,0]) \in \Delta(T)$ , and  $\Delta(T)$  is also isomorphic to  $T$ , we conclude that  $([n,0],0) \in L(S)$  implies that  $n = 0$ . A similar argument works for the points of the form  $([m,r],[n,0])$  with  $n \leq m$ , and we conclude that  $L(S) = \{([n,0],1), (1,[n,0]) : n \in \mathbb{N}\} \cup \{(1,1), (0,0)\}$ . Fix  $m \in \mathbb{N}$ . Then, for all  $n \in \mathbb{N}$ ,  $([n,0],1)(1,[m,0]) = (0,0)$  (this product is in  $L(S)$ ). Hence, since  $(1,1) = \sup_n ([n,0],1)$ , it follows that  $L(S)$  is not lower continuous, and so  $L(S)$  cannot have a compact semilattice topology.

Question: Let  $S \in \underline{CS}$ , and suppose that  $L(S) \in \underline{CS}$ . Is then  $L(S) \in \underline{CL}$ ?