

NAME: Hans Dobbertin

Date: 

M	D	Y
11	18	82

TOPIC: Distributive semilattices, Heyting algebras and V-homomorphisms

## REFERENCES:

- [C] Compendium.  
 [D1] H. Dobbertin, On Vaught's criterion for isomorphisms of countable Boolean algebras, Algebra Univ. 15 (1982), in press.  
 [D2] H. Dobbertin, Verfeinerungsmonoide, Vaught Monoide und Boolesche Algebren, Dissertation, Universität Hannover, 1982.  
 [G] G. Grätzer, Lattice Theory: First Concepts and Distributive Lattices, W. H. Freeman, San Francisco, 1971.  
 [K] J. Katonien, The structure of countable Boolean algebras, Ann. Of Math. 108 (1978), 41-89.  
 [M] D. Myers, Structures and measures on Boolean algebras, unpublished manuscript.

The study of the monoid, under direct sums, of all isomorphism

types of countable Boolean algebras has led to the notion of a refinement monoid  $[M, D]$ , cf. [K]:

A commutative monoid  $M = (M; +, 0)$  is called a refinement monoid provided that

(RM1)  $x + y = 0$  only for  $x = y = 0$  ( $x, y \in M$ ),

(RM2)  $M$  has the refinement property, that is, whenever  $\sum x_i = \sum y_j$

for  $x_i, y_j \in M$  ( $i < n, j < m$ ) then there are  $z_{ij} \in M$

with  $x_i = \sum_j z_{ij}$  and  $y_j = \sum_i z_{ij}$ .

A homomorphism  $h : M \rightarrow N$  between commutative monoids is said to be a V-homomorphism if  $h(x) = y_1 + y_2$  ( $x \in M, y_1, y_2 \in N$ ) implies

$x = x_1 + x_2$  and  $h(x_i) = y_i$  for some  $x_1, x_2 \in M$ , and  $h(x) = 0$

only for  $x = 0$  ( $x \in M$ ). Observe that a V-homomorphic image of a refinement monoid is again a refinement monoid.

PROPOSITION 1. A semilattice  $L = (L; +, 0)$  with zero is distributive (in the sense of [G; p. 117]) iff  $L$  is a refinement monoid.

It is well-known that the category of distributive semilattices with zero and homomorphisms having the property that pre-images of prime filters are always prime filters is dually equivalent to the category of Stone spaces (sober  $T_0$ -spaces having a base of compact sets) and strongly continuous mappings (pre-images of compact-open sets are compact-open); see [G; 2.11].

Let  $DSL$  be the category of distributive semilattices with zero and V-homomorphisms. We want to supply the category  $STS$  of Stone

spaces with suitable morphisms so that  $STS$  and  $DSL$  become equivalent categories. First let us call a subset  $U$  of a space  $X$  almost-open if there is a smallest open set, say  $\tilde{U}$ , containing  $U$ , and  $U$  is a strict subset of  $\tilde{U}$  (i. e., the inclusion map from  $U$  into  $\tilde{U}$  is strict [C; V.5.8]). Note that, for instance, every space is almost-open in its sobrification. Of course, open sets are almost-open. Now suppose that  $X$  and  $Y$  are Stone spaces, then  $\text{mor}(X, Y)$  consists of all continuous functions from  $X$  into  $Y$  mapping open sets (or equivalently, almost-open sets) onto almost-open sets. Thus all continuous-open mappings are morphisms of  $STS$ . Probably, the converse is false. However, I have no counterexample.

We call a mapping  $h : L \rightarrow K$  between complete lattices a strong V-homomorphism if  $h$  is Sup-preserving,  $h(x) = \sup_{i \in I} y_i$  always implies  $x = \sup_{i \in I} x_i$  and  $h(x_i) = y_i$  for some elements  $x_i$ , and  $h(x) = 0$  only for  $x = 0$ .

LEMMA 2. Let  $H_1$  and  $H_2$  be complete Heyting algebras, and suppose that  $g : H_1 \rightarrow H_2$  and  $h : H_2 \rightarrow H_1$  form an adjunction [C; p. 18]. Then the following are equivalent:

- (i)  $g$  preserves Sups, Infs and  $\Rightarrow$ ,  
 (ii)  $h$  is a strong V-homomorphism.

Let  $AHA_0$  (resp.  $AHA_j$ ) be the category of algebraic complete Heyting algebras and strong V-homomorphisms (resp.  $\Rightarrow$ -preserving, complete homomorphisms).

PROPOSITION 3. The categories  $STS, DSL, AHA_0, AHA_j^{op}$  are equivalent.

Of course, the emphasis in Proposition 3 lies on the morphisms; on the object level this is well-known.

THEOREM 4. [D2] Let  $L$  be a distributive semilattice with zero. If  $L$  is a lattice or  $|L| \leq \aleph_1$  then  $L$  is a V-homomorphic image of some generalized Boolean lattice.

Question A. Does Theorem 4 hold for all  $L$ ?

A "dual version" of Thm. 4 is the following: Let  $H$  be an algebraic complete Heyting algebra such that the set  $K(H)$  of compact

elements of  $H$  is a lattice or  $|K(H)| \leq \aleph_1$ , then  $H$  is embeddable into the ideal lattice  $\text{Id}(B)$  of some generalized Boolean algebra  $B$  under a mapping preserving sups, infs and  $\Rightarrow$ . As a consequence, every Heyting algebra can be embedded into  $\text{Id}(B)$  for some Boolean algebra  $B$  (under a mapping preserving sups, infs and  $\Rightarrow$ ). A similar result for distributive pseudo-complemented lattices has been shown by Lakser (see [G; p. 180]).

Question B. Is every Stone space  $X$  the image of a locally compact zero-dimensional Hausdorff space under a continuous-open mapping?

It is not difficult to see that if  $X$  is first-countable then, for all Stone spaces  $Y$ ,  $\text{mor}(Y, X)$  consists only of continuous-open mappings. Thus, in this case, it follows from Thm. 4 that Question B has an affirmative answer provided that the set  $L(X)$  of compact-open subsets of  $X$  is closed under finite intersections or  $|L(X)| \leq \aleph_1$ .

*Erwas erweitere Fassung  
des "Meas" vom 12.11.82*

*Gruf  
HD*